

**IN PRESS, BRITISH JOURNAL OF PSYCHIATRY**

**Word count:** Abstract: 150

Text: 3,999

Tables: 3

**Childhood Adversities and Post-traumatic Stress Disorder: Evidence for Stress**

**Sensitization in the World Mental Health Surveys\***

Running title: Childhood adversities and PTSD

Katie A. McLaughlin, Karestan C. Koenen, Evelyn J. Bromet, Elie G. Karam, Howard Liu, Maria Petukhova, Ayelet Meron Ruscio, Nancy A. Sampson, Dan J. Stein, Sergio Aguilar-Gaxiola, Jordi Alonso, Guilherme Borges, Koen Demyttenaere, Rummyana V. Dinolova, Finola Ferry, Silvia Florescu, Giovanni de Girolamo, Oye Gureje, Norito Kawakami, Sing Lee, Fernando Navarro-Mateu, Marina Piazza, Beth-Elle Pennell, José Posada-Villa, Margreet ten Have, Maria Carmen Viana, Ronald C. Kessler

\*The paper is submitted on behalf of the World Health Organization World Mental Health Survey Collaborators – see the Appendix for details.

May 2017

**Author Affiliations**

**Katie A. McLaughlin**, PhD, Department of Psychology, University of Washington, Seattle, Washington, USA; **Karestan C. Koenen**, PhD, Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA; **Evelyn J. Bromet**, PhD, Department of Psychiatry, Stony Brook University School of Medicine, Stony Brook, New York, USA; **Elie G. Karam**, MD, Department of Psychiatry and Clinical Psychology, Faculty of Medicine, Balamand University, Beirut, Lebanon; Department of Psychiatry and Clinical Psychology, St George Hospital University Medical Center, Beirut, Lebanon; Institute for Development Research Advocacy and Applied Care (IDRAAC), Beirut, Lebanon; **Howard Liu**, SM, Department of Health Care Policy, Harvard Medical School, Boston, Massachusetts, USA; **Maria Petukhova**, PhD, Department of Health Care Policy, Harvard Medical School, Boston, Massachusetts, USA; **Ayelet Meron Ruscio**, PhD, Department of Psychology, University of Pennsylvania, Philadelphia, Pennsylvania, USA; **Nancy A. Sampson**, BA, Department of Health Care Policy, Harvard Medical School, Boston, Massachusetts, USA; **Dan J. Stein**, FRCPC, PhD, Department of Psychiatry and Mental Health, University of Cape Town, Cape Town, Republic of South Africa; **Sergio Aguilar-Gaxiola**, MD, PhD, Center for Reducing Health Disparities, UC Davis Health System, Sacramento, California, USA; **Jordi Alonso**, MD, DrPH, IMIM-Hospital del Mar Research Institute, Parc de Salut Mar; Pompeu Fabra University (UPF); CIBER en

Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain; **Guilherme Borges**, DrSc, Instituto Nacional de Psiquiatría, Calzada Mexico Xochimilco No 101- Col. San Lorenzo Huipulco, Mexico; **Koen Demyttenaere**, MD, PhD, Department of Psychiatry, University Hospital Gasthuisberg, Katholieke Universiteit Leuven, Leuven, Belgium; **Rumyana V. Dinolova**, MD, National Center of Public Health and Analyses, Sofia, Bulgaria; **Finola Ferry**, PhD, Bamford Centre for Mental Health and Wellbeing, Ulster University, Coleraine, United Kingdom; **Silvia Florescu**, MD, PhD, National School of Public Health, Management and Development, Bucharest, Romania; **Giovanni de Girolamo**, MD, Unit of Epidemiological and Evaluation Psychiatry, IRCCS -St. John of God Clinical Research Centre, Brescia, Italy; **Oye Gureje**, MD, PhD, FRCPsych, Department of Psychiatry, University College Hospital, Ibadan, Nigeria; **Norito Kawakami**, MD, DMSc, Department of Mental Health, School of Public Health, The University of Tokyo, Tokyo, Japan; **Sing Lee**, MB, BS, Department of Psychiatry, Chinese University of Hong Kong, Tai Po, Hong Kong; **Fernando Navarro-Mateu**, MD, PhD, UDIF-SM, Subdirección General de Planificación, Innovación y Cronicidad, Servicio Murciano de Salud; IMIB-Arrixaca; CIBERESP-Murcia, Murcia, Spain; **Beth-Ellen Pennell**, MA, Survey Research Center, Institute for Social Research, University of Michigan, Ann Arbor, Michigan, USA; **Marina Piazza**, ScD, MPH, Universidad Cayetano Heredia; National Institute of Health, Lima, Peru ; **José Posada-Villa**, MD, Faculty of Social Sciences Colegio Mayor de Cundinamarca University, , Bogota, Colombia; **Margreet ten Have**, PhD, Trimbos-Instituut, Netherlands Institute of Mental Health and Addiction, Utrecht, Netherlands; **Maria Carmen Viana**, MD, PhD, Department of Social Medicine, Federal University of Espírito Santo, Vitoria, Brazil; **Ronald C. Kessler**, PhD, Department of Health Care Policy, Harvard Medical School, Boston, Massachusetts, USA

**Correspondence:** Katie A. McLaughlin, Department of Psychology, University of Washington, Box 351525, Seattle, WA, 98195; Phone 206-616-7863; [mclaughk@uw.edu](mailto:mclaughk@uw.edu)

## ABSTRACT

**Background:** Although childhood adversities (CAs) are known to predict increased risk of post-traumatic stress disorder (PTSD) after traumatic experiences (TEs), it is unclear whether this association varies by CA or TE types or by age.

**Aims:** To examine variation in associations of CAs with PTSD according to CA types, TE types, and life-course stage.

**Method:** Epidemiological data were analyzed from the World Mental Health Surveys (N=27,017).

**Results:** Four CAs (physical and sexual abuse, neglect, parent psychopathology) were associated with similarly increased odds of PTSD following TEs (OR=1.8), while the other 8 CAs assessed did not predict PTSD. CA-PTSD associations did not vary across TE types, but were stronger in childhood-adolescence and early-middle adulthood than later adulthood.

**Conclusion:** CAs are differentially associated with PTSD, with the strongest associations in childhood-adolescence and early-middle adulthood. Consistency of associations across TE types suggests that CAs are associated with generalized vulnerability to PTSD following TEs.

## INTRODUCTION

Child maltreatment has repeatedly been shown to predict post-traumatic stress disorder (PTSD) among adults exposed to traumatic events (TEs)(1-5). It is unknown whether or not a broader set of childhood adversities (CAs) are similarly associated with increased vulnerability to PTSD following TEs(6), although the associations of more general stressful life events with anxiety, mood, and substance disorders are known to be elevated among individuals who have experienced a wide range of CAs(6-12). Prior work has documented that CAs reflecting maladaptive family functioning (MFF)—including child maltreatment, parent psychopathology, and family violence—are more strongly associated with onset of mental disorders than other CAs and that the joint associations of MFF CAs with onset of mental disorders are sub-additive (i.e., the incremental associations of additional CAs decrease as the number of CAs increases) (13-16). Comparable analyses have not been carried out, though, with respect to the associations of CAs with subsequent onset of PTSD in the wake of TE exposure. It is also unclear whether CAs are associated with a generalized vulnerability to PTSD following any TE type or whether these associations are more pronounced for specific types of TEs. With regard to the latter possibility, it is known that PTSD risk varies markedly across TE types(17-19) and that prior exposure to some TE types, most notably those involving physical violence victimization, are associated with elevated PTSD risk following subsequent TEs of the same type.(19) It is plausible in light of this evidence to think that the associations of CAs with PTSD might vary across TE types. Finally, although CAs have been shown to predict onset of anxiety, mood, and substance disorders similarly at every stage of the life-span(13-16), prior work has not examined whether vulnerability to PTSD among those with a history of CAs varies according to life-course stage.

In the current study, we analyze the general population epidemiological data in the WHO World Mental Health (WMH) Surveys, a cross-national series of community epidemiological surveys of the prevalence and correlates of common mental disorders, to examine whether CAs are associated with increased risk of PTSD after TE exposure. The large sample size of the WMH surveys allows us to consider variation in the associations of CAs with PTSD as a function of type of CA, type of TE, and life-course stage.

## METHODS

### Samples

Data come from the 20 WMH surveys in 18 countries that assessed CAs and used an expanded assessment of PTSD (described below) to examine PTSD associated with a *randomly-selected* TE. (Appendix Table 1) These surveys included 10 in countries classified by the World Bank(20) as high-income (national surveys in Belgium, France, Germany, Italy, the Netherlands, Northern Ireland, Spain and the United States along with a survey of metropolitan areas in Japan and the Murcia region of Spain), 7 in countries classified as upper-middle-income (national surveys in Bulgaria, Lebanon, Romania and South Africa along with a survey of all non-rural areas in Mexico and regional surveys in Sao Paulo Brazil and Medellin Colombia), and 3 in countries classified as lower-middle/low income (national surveys in Peru and Ukraine and a survey of all non-rural areas in Colombia). The two Colombia surveys were classified as being in different country-income groups because the World Bank ranking of Colombia's income-level changed between the times of the two surveys. All surveys were based on multi-stage clustered area probability samples of adult household residents. Response rates ranged from 45.9% (France) to 97.2% (Colombia) and had a weighted mean of 70.9% across surveys. A detailed description of sampling procedures is presented elsewhere(21).

## Field procedures

Interviews were administered face-to-face in respondent's homes after obtaining informed consent using procedures approved by local Institutional Review Boards. The interview schedule was developed in English and translated into other languages using a standardized WHO translation, back-translation, and harmonization protocol.(22) Training and field procedures were supervised by the WMH Data Collection Coordination Centre to guarantee cross-national consistency(22).

Interviews were administered in two parts. Part I, administered to all respondents, assessed core DSM-IV mental disorders (n=83,805). Part II was administered to all respondents who met lifetime criteria for any Part I disorder and a probability subsample of other Part I respondents (n=42,430). Part II assessed additional disorders and correlates. Questions about TEs and PTSD were included in Part II. Part II respondents were weighted to adjust for differential probabilities of selection and deviations between the sample and population demographic-geographic distributions. Details about WMH weighting are presented elsewhere(21). The subsample of Part II respondents who reported lifetime exposure to one or more TEs (n=27,017) is the focus of the current report.

## Measures

**Exposure to traumatic experiences:** Part II respondents were asked about lifetime exposure to 27 different types of TEs in addition to two open-ended questions about exposure to “*any other*” TE and to any “*private*” TE the respondent did not want to discuss. Positive responses were followed by probes to assess number of lifetime exposures and age at first exposure to each TE type. Exploratory factor analysis in the WMH sample found 6 TE groups:(23) 5 reflecting exposure to organized violence (e.g., civilian in a war zone, relief

worker in a war zone, refugee); 5 related to participation in organized violence (e.g., combat experience, witnessed atrocities); 3 reflecting physical violence victimization (witnessed violence at home as a child; beaten by a caregiver as a child; victim of assault); 7 related to sexual violence (e.g., raped, sexually assaulted, beaten by a romantic partner); 6 involving accidents/injuries (e.g., natural disaster, toxic chemical exposure, motor vehicle accident); and 3 that were not strongly correlated with any other TEs (mugged or threatened with a weapon, manmade disaster other than chemical exposure, unexpected death of a loved one).

**PTSD:** Mental disorders were assessed with the Composite International Diagnostic Interview (CIDI)(24), a fully-structured interview administered by trained lay interviewers that assesses DSM-IV disorders. PTSD was assessed in relation to the lifetime TE the respondent identified as “worst” (i.e., as causing the most severe-persistent core symptoms of PTSD) and in response to one randomly-selected TE out of all TEs the respondent reported experiencing. The random TE could be the same as the TE selected as the worst. Population estimates of PTSD prevalence and conditional risk of PTSD are likely over-estimated when assessed in relation to a respondent’s worst TE.(17, 18) Consequently, we focus here on PTSD associated with randomly-selected TEs, an approach that has previously been used in cross-national research from the WMH Surveys.(19, 25) To do so, we created a population-level representative sample of TEs. As each respondent had only one randomly-selected TE no matter how many lifetime TEs they experienced, TEs that occurred to respondents who experienced a high number of TEs were under-represented. We corrected for this by weighting the randomly-selected TE by the inverse of its probability of selection and then multiplying this weight by the Part II weight to generate a sample representative of all TEs experienced by all respondents. These weighted data no longer represent the population of *respondents* but the population of *all TEs* experienced in

the population. The consolidated weight was standardized in each survey to equal the number of respondents assessed for randomly-selected TEs.

As detailed elsewhere(26), blinded clinical reappraisal interviews with the Structured Clinical Interview for DSM-IV (SCID) conducted in four WMH countries found CIDI-SCID concordance for DSM-IV PTSD to be moderate (AUC=.69). Sensitivity and specificity were .38 and .99, respectively, resulting in a likelihood ratio positive (LR+) of 42.0, which is well above the threshold of 10 typically used to consider screening scale diagnoses definitive.(27) Consistent with the high LR+, the proportion of CIDI cases confirmed by the SCID was 86.1%. This means that the vast majority of CIDI/DSM-IV PTSD cases would be diagnosed with PTSD by a trained clinician.

**Childhood adversities:** Twelve CAs occurring before age 18 years were assessed. These included 3 types of interpersonal loss (parental death, parental divorce, and other loss of contact with parents/caregivers), 4 types of parental maladjustment (psychopathology, substance abuse, criminality, and family violence), 3 types of maltreatment (physical abuse, sexual abuse, neglect), respondent serious physical illness, and economic adversity. These CAs align with the Adverse Childhood Experiences Study,(28) although we did not assess emotional abuse and additionally assessed economic adversity and serious physical illness.

Physical abuse of the respondent by caregivers was assessed with a modified version of the Conflict Tactics Scale (CTS)(29) and with an item from the TE section of the CIDI. Sexual abuse was assessed with questions from the CIDI regarding sexual assault, attempted rape, and rape. Neglect was assessed with questions that assessed frequency of not having adequate food, clothing, or medical care, having inadequate supervision, and being required to do chores that were age-inappropriate.(30)



Parental criminality assessed whether a parent engaged in criminal activities or was ever arrested or sent to prison. Parent psychopathology (major depression, generalized anxiety disorder, panic disorder, suicide attempt) and substance abuse were assessed with a revised version of the Family History Research Diagnostic Criteria Interview.(31, 32) Family violence was assessed with the modified CTS and an item in the TE section of the CIDI. Economic adversity was assessed with questions about whether the respondent's family received welfare or other government assistance or often did not have enough money to pay for basic necessities. Physical illness was assessed with a standard chronic conditions checklist.

Four of these CAs are also TEs: sexual abuse; physical abuse; witnessing family violence during childhood; and childhood physical illness. If the randomly-selected TE was one of these four CAs, it was not included among the CA measures. Several WMH countries omitted some CAs (neglect in South Africa; parental divorce and neglect in the six Western European countries). Rather than exclude these countries or exclude the missing CAs from analysis, we imputed individual-level missing values using multiple imputation.(33)

Prior factor analysis in this sample identified a cluster of highly-correlated CAs reflecting maladaptive family functioning (MFF) that included parent psychopathology, parent substance abuse, parent criminality, family violence, physical abuse, sexual abuse, and neglect.(13) We analyzed these MFF CAs separately from Other CAs (parental death and divorce, other parental loss, physical illness, economic adversity), which exhibit weaker inter-correlations.

### **Analysis Methods**

Cross-tabulations were used in this weighted-standardized dataset to estimate prevalence of exposure to CAs associated with randomly-selected TEs. Associations of these CAs with PTSD in the full sample of randomly-selected TEs were then estimated using logistic regression

models controlling sex, age, TE type, and history of prior TEs of the same type as the randomly-selected TE for TEs where our previous work shows that prior exposure significantly predicts PTSD risk.(19) Dummy predictor variables were included for survey to adjust for between-survey differences in aggregate PTSD prevalence. A series of multivariate models was then estimated using a modeling approach we have employed previously(13, 15, 16) that distinguishes between MFF and Other CAs and evaluates separate and joint associations of multivariate CA profiles with PTSD onset after TE exposure. We began with a series of bivariate models (M1) examining each CA separately. We then estimated an additive multivariate model that included a separate predictor variable for each of the 12 CAs (M2) followed by a model that included predictor variables for number of CAs without variables for specific CA types (M3). The latter model included one variable for the number of MFF CAs (range: 0-4) and two dummy variables indicating exposure to exactly 1 or 2+ Other CAs. Only MFF CAs that were significant in the bivariate models were included in the 0-4 count. We then estimated a model that included dummy variables for CA types in addition to variables for number of MFF and Other CAs (M4). The next model deleted the 4 MFF CA type variables that were included in the MFF CA count variable but retained the other type variables (M5). The next model (M6) included the same CA type variables as M5 but included dummy variables for exact numbers of MFF CAs rather than the count variable. A final model (M7) included only the MFF CA count variable and dropped all other CA variables.

Inspection of mean-squared error (MSE) and area under the receiver operating characteristic curve (AUC) based on replicated 10-fold cross-validation with 10 replications was used to select the best multivariate model among M2-M7 to predict PTSD in response to the randomly-selected TE. Additional models examined whether associations of CAs with PTSD

varied by TE type and timing of TE exposure with interaction terms added to the best-fitting model. Composite TE measures used in these interactions were based on prior analysis showing that odds of PTSD in response to the randomly-selected TE were equivalent for all but 8 TE types.(19) Here, we included predictor variables for those 8 TE types and one variable for all other TE types. Given that all respondents were exposed to one randomly-selected TE, the indeterminacy of the solution created by including an odds-ratio (OR) for each TE type was resolved by scaling the product of all TE-specific ORs to 1.0. Any TE-specific ORs significantly different from 1.0 can be interpreted as having significantly higher- or lower-than-average odds of PTSD; each TE type was given equal weight in computing this average. Interactions between CAs and age of exposure to the randomly-selected TE classified age into 5 groups (0-12, 13-24, 25-44, 45-59, and 60+ years).

Logistic regression coefficients and standard errors were exponentiated and are reported as ORs with 95% confidence intervals (CIs). Statistical significance was evaluated using .05-level two-sided tests based on the design-based Taylor series method(34) implemented in the SAS software system(35) to adjust for the weighting and clustering of observations.

## RESULTS

### CA Prevalence

Prevalence estimates of CAs associated with representative TEs are shown in Table 1. The proportions of TEs occurring to individuals with a history of CAs range from 3.8% (sexual abuse) to 20.9% (physical abuse). It is noteworthy that all prevalence estimates in the *sample of TEs* are higher than in the *sample of respondents*, indicating that CAs are positively associated with number of TEs experienced. For example, 10.7% of respondents reported exposure to physical abuse, but 20.9% of all TEs occurred to individuals exposed to physical abuse. The

most elevated CA prevalence estimates in the TE sample relative to the sample of respondents involve the MFF CAs of family violence, physical abuse, and sexual abuse.

**(Table 1 here)**

### **Associations of childhood adversities with PTSD**

In bivariate models (M1) that considered one CA at a time, only MFF CAs were significant predictors of PTSD after the randomly-selected TE, with ORs in the range of 1.6-2.2. (Table 2) MFF CAs associated with PTSD included physical and sexual abuse, neglect, and parent psychopathology. The ORs for incremental associations of these 4 MFF CAs with PTSD were lower in the additive multivariate model (M2) that included all CA types. Three of the four MFF CAs remained significant in that model (parent psychopathology, sexual abuse, neglect), with ORs in the range of 1.8-2.0. MFF CAs, as a set, were significantly associated with PTSD in M2 ( $\chi^2_7=40.8$ ,  $p<.001$ ), but Other CAs were not ( $\chi^2_5=5.8$ ,  $p=.33$ ). In addition, the strength of the ORs varied significantly across the different types of MFF CAs ( $\chi^2_6=18.5$ ,  $p=.005$ ) but not Other CAs ( $\chi^2_4=5.8$ ,  $p=.22$ ).

**(Table 2 here)**

In multivariate model M3, which considered only number and not type of CAs, the number of MFF CAs was significantly associated with PTSD (OR=1.7,  $p<.001$ ), but the number of Other CAs was not ( $\chi^2_2=0.4$ ,  $p=.84$ ). (Table 2)

Subsequent models (M4-M6) included information about both type and number of CAs. We examined models including variables for number of MFF and Other CAs as well as type variables for all CAs (M4), and type variables for CAs excluding the 4 MFF CAs included in the count variable (M5). We also examined a model that included variables for exactly 1, 2, 3, and 4 MFF CAs rather than the 0-4 count (M6). However, the best-fitting model (M7) included only

the count variable for the 4 individually-significant MFF CAs (Table 2). Consistent with M7 being the best model, neither the other MFF CAs not included in the count variable nor Other CAs were associated with PTSD in M5 or M6, indicating an absence of meaningful associations of CAs other than physical and sexual abuse, neglect, and parent psychopathology with PTSD. (Table 2) In the best-fitting model (M7), number of MFF CAs was significantly associated with PTSD (OR=1.8,  $p < .001$ ).

### **Differential associations by TE type and timing of TE exposure**

To determine whether the association of number of MFF CAs with PTSD varied as a function of the type of TE, we added interactions between this variable and 9 indicators for TE type to the best-fitting CA model (M7). The association of number of MFF CAs with PTSD did not vary across TE types ( $\chi^2_8=13.9$ ,  $p=.08$ ). The ORs for the interaction term were in the range of 0.8-1.6 with the exception of natural disaster (OR=0.3), witnessing atrocities (OR=0.4), and being kidnapped (OR=3.1). (Table 3)

#### **(Table 3 here)**

Next, we examined interactions between number of MFF CAs and timing of TE exposure. Here, the association of number of MFF CAs with PTSD varied depending on age of TE exposure ( $\chi^2_4=16.4$ ,  $p=.002$ ). Specifically, coefficients were positive and associations were significant for TEs occurring during childhood (OR=1.6,  $p=.007$ ), adolescence (OR=1.8,  $p=.001$ ), and adulthood through age 44 (OR=1.9,  $p < .001$ ) but not for TEs occurring in later-middle age (ages 45-59; OR=1.5,  $p=.16$ ), and the OR was reversed among respondents aged 60+ (OR=0.1,  $p < .001$ ). (Table 3)

## **DISCUSSION**

We found that exposure to some, but not all, types of CAs is associated with increased likelihood of developing PTSD. Of the wide range of CAs examined, only physical and sexual abuse, neglect, and parent psychopathology were associated with elevated risk of developing PTSD following a TE. These findings extend prior work documenting higher prevalence of past-year PTSD following a past-year TE among individuals with high levels of CA exposure(6) and greater risk of developing PTSD among individuals who were maltreated as children(1-5) by documenting parental psychopathology as a CA associated with PTSD in addition to maltreatment and by documenting that these vulnerabilities do not extend to the other CA types considered here. Our finding of an association of parental psychopathology with PTSD is broadly consistent with evidence that parent psychopathology has robust but largely non-specific associations with offspring psychopathology.(36, 37) Unlike prior work suggesting that the associations of CAs with mental disorders are sub-additive, such that the incremental effects of additional CAs get relatively smaller as the number of CAs increases,(14-16) our results indicate that the associations of CAs involving maltreatment and parent psychopathology with PTSD are additive. That is, each additional exposure to these specific CAs is associated with increased odds of developing PTSD following a TE that is relatively consistent as the number of CAs increases.

The association of CAs involving child maltreatment and parent psychopathology with PTSD did not vary across a wide range of TE types. This is surprising, given extensive previous research showing that conditional risk of PTSD varies considerably across TE types.(17-19) Prior work on stress sensitization has produced conflicting findings in terms of whether vulnerability to stressors among individuals with a history of CAs is general or specific to certain types of stressors. Some studies have shown that relatively mild stressors are more likely to

trigger depression among those with a history of CA than those without,(10, 38) whereas other work indicates that CA exposure magnifies the association between relatively severe stressors and psychopathology.(6) Our failure to document variation in the associations of CAs with PTSD as a function of TE type is consistent with the possibility that CAs produce a generalized vulnerability to PTSD following TEs.

The associations of CAs and PTSD varied by life-course stage, with the associations observable during childhood, adolescence, and early-middle adulthood, but not during late-middle age or later adulthood. We are unaware of prior research examining this specification, though it is noteworthy that most prior studies documenting an interaction between CAs and later stressors in predicting mental disorders focused on adolescence or young adulthood.(10-12, 38, 39) There are two possible interpretations of this finding. The first is that the association of CAs with PTSD wanes later in life due to increased temporal distance from the CA, variations in the features of TEs across the lifespan, or protective factors that increase in later life. Alternatively, it might reflect recall biases in CA reports that are strongest among older adults, where the recall interval is longest. PTSD is associated with over-general autobiographical memory,(40) and it is possible that this effect is more pronounced in older adults. Under-reporting of CAs would attenuate CA-PTSD associations in the oldest group if PTSD symptoms are not under-reported. The fact that CAs were negatively associated with PTSD among older adults might be due to selection of especially resilient individuals with CA histories into the later years of life.

Why might exposure to child maltreatment and parent psychopathology influence vulnerability to PTSD? And why were only these specific MFF CAs associated with PTSD? One potential mechanism involves heightened salience of threat cues and magnified emotional reactions to potential threats. Child maltreatment is associated with elevated emotional reactivity

to negative stimuli, as measured using self-report,(41, 42) ecological momentary assessments of daily responses to stressful life events,(43) and amygdala response to negative stimuli.(44-47) Similar patterns have been observed in children of parents with psychopathology.(48, 49) Heightened amygdala reactivity to threat may be a key pathway through which maltreatment and parent psychopathology influence later risk for PTSD. Prospective studies in both adolescents and military samples have shown that elevated amygdala response to threat cues assessed *prior* to trauma exposure predicts the subsequent onset of PTSD symptoms following a TE.(50, 51) Amygdala reactivity to threat cues decreases in older age,(52) which may have contributed to the lack of CA-PTSD associations in older adults, although this does not account for the negative CA-PTSD association among older respondents. The specificity of this pattern of neural response to CAs involving direct exposure to threat(53) or parent psychopathology(46, 47) may explain, in part, why CAs involving interpersonal loss and economic adversity were not associated with PTSD vulnerability. Other mechanisms are also likely to play a role in the associations of these specific MFF CAs with PTSD, including habitual use of maladaptive emotion regulation strategies (e.g., rumination) and low social support, each of which is associated with child maltreatment and parent psychopathology as well as risk for PTSD.(1, 41, 54-56)

This study is limited by retrospective reports of CAs, trauma exposure, and PTSD symptoms. Recall bias of CAs primarily involves under-reporting,(57-59) which likely attenuated associations of CAs with PTSD. The absence of CA associations with PTSD in the oldest respondents could reflect under-reporting that was most pronounced among respondents with the longest recall periods.(15) Additionally, assessment of PTSD with a fully-structured diagnostic interview rather than clinician-administered interview likely introduced imprecision.



The CIDI diagnoses were relatively conservative, meaning that low-severity cases of PTSD may have been missed. We did not assess all relevant CAs, most notably emotional abuse.

Determining whether emotional abuse influences vulnerability to PTSD in a similar manner as physical and sexual abuse is an important direction for future work. Finally, our measure of economic adversity included assessment of receipt of governmental assistance, which varies widely across countries in terms availability and eligibility criteria and may underestimate economic adversity in some countries.

Despite these limitations, we replicate prior work indicating that child maltreatment is associated with heightened vulnerability to PTSD following TEs occurring later in development. We extend earlier findings by showing that a similar association exists with regard to parent psychopathology but not other forms of CA, that the association of CAs involving child maltreatment and parent psychopathology with PTSD are additive and do not vary across TE types, and that CA-PTSD associations are limited to childhood, adolescence, and early-middle adulthood. These results build on a growing body of evidence indicating that early experiences of adversity increase vulnerability to psychopathology following stressors later in development and suggest that CAs involving child maltreatment and parental psychopathology are associated with heightened risk for PTSD following a TE.

## ACKNOWLEDGMENTS

The paper is submitted on behalf of the World Health Organization World Mental Health Survey collaborators: Sergio Aguilar-Gaxiola, MD, PhD, Ali Al-Hamzawi, MD, Mohammed Salih Al-Kaisy, MD, Jordi Alonso, MD, DrPH, Laura Helena Andrade, MD, PhD, Corina Benjet, PhD, Guilherme Borges, ScD, Evelyn J. Bromet, PhD, Ronny Bruffaerts, PhD, Brendan Bunting, PhD, Jose Miguel Caldas de Almeida, MD, PhD, Graca Cardoso, MD, PhD, Somnath Chatterji, MD, Alfredo H. Cia, MD, Louisa Degenhardt, PhD, Koen Demeyttenaere, MD, PhD, John Fayyad, MD, Silvia Florescu, MD, PhD, Giovanni de Girolamo, MD, Oye Gureje, PhD, DSc, FRCPsych, Josep Maria Haro, MD, PhD, Yanling He, MD, Hristo Hinkov, MD, Chi-yi Hu, PhD, MD, Yueqin Huang, MD, MPH, Peter de Jonge, PhD, Aimee Nasser Karam, PhD, Elie G. Karam, MD, Norito Kawakami, MD, DMSc, Ronald C. Kessler, PhD, Andrzej Kiejna, MD, PhD, Viviane Kovess-Masfety, MD, PhD, Sing Lee, MB, BS, Jean-Pierre Lepine, MD, Daphna Levinson, PhD, John McGrath, MD, PhD, Maria Elena Medina-Mora, PhD, Jacek Moskalewicz, DrPH, Fernando Navarro-Mateu, MD, PhD, Beth-Ellen Pennell, MA, Marina Piazza, MPH, ScD, Jose Posada-Villa, MD, Kate M. Scott, PhD, Tim Slade, PhD, Juan Carlos Stagnaro, MD, PhD, Dan J. Stein, FRCPC, PhD, Margreet ten Have, PhD, Yolanda Torres, MPH, Dra.HC, Maria Carmen Viana, MD, PhD, Harvey Whiteford, PhD, David R. Williams, MPH, PhD, Bogdan Wojtyniak, ScD.

## FINANCIAL SUPPORT

The World Health Organization World Mental Health (WMH) Survey Initiative is supported by the National Institute of Mental Health (NIMH; R01 MH070884 and R01 MH093612-01), the John D. and Catherine T. MacArthur Foundation, the Pfizer Foundation, the US Public Health Service (R13-MH066849, R01-MH069864, and R01 DA016558), the Fogarty International Center (FIRCA R03-TW006481), the Pan American Health Organization, Eli Lilly and Company, Ortho-McNeil Pharmaceutical, Inc., GlaxoSmithKline, and Bristol-Myers Squibb. We thank the staff of the WMH Data Collection and Data Analysis Coordination Centres for assistance with instrumentation, fieldwork, and consultation on data analysis. RK and KCK were supported by the US National Institute of Mental Health (NIMH) under Award Number R01MH101227.

The São Paulo Megacity Mental Health Survey is supported by the State of São Paulo Research Foundation (FAPESP) Thematic Project Grant 03/00204-3. The Bulgarian Epidemiological Study of common mental disorders EPIBUL is supported by the Ministry of Health and the National Center for Public Health Protection. The Colombian National Study of Mental Health (NSMH) is supported by the Ministry of Social Protection. The Mental Health Study Medellín – Colombia was carried out and supported jointly by the Center for Excellence on Research in Mental Health (CES University) and the Secretary of Health of Medellín. The ESEMeD project is funded by the European Commission (Contracts QLGS-1999-01042; SANCO 2004123, and EAHC 20081308), (the Piedmont Region (Italy)), Fondo de Investigación Sanitaria, Instituto de Salud Carlos III, Spain (FIS 00/0028), Ministerio de Ciencia y Tecnología, Spain (SAF 2000-158-CE), Departament de Salut, Generalitat de Catalunya, Spain, Instituto de Salud Carlos III (CIBER CB06/02/0046, RETICS RD06/0011 REM-TAP), and other local

agencies and by an unrestricted educational grant from GlaxoSmithKline. The World Mental Health Japan (WMHJ) Survey is supported by the Grant for Research on Psychiatric and Neurological Diseases and Mental Health (H13-SHOGAI-023, H14-TOKUBETSU-026, H16-KOKORO-013, H25-SEISHIN-IPPAN-006) from the Japan Ministry of Health, Labour and Welfare. The Lebanese Evaluation of the Burden of Ailments and Needs Of the Nation (L.E.B.A.N.O.N.) is supported by the Lebanese Ministry of Public Health, the WHO (Lebanon), National Institute of Health / Fogarty International Center (R03 TW006481-01), anonymous private donations to IDRAAC, Lebanon, and unrestricted grants from, Algorithm, AstraZeneca, Benta, Bella Pharma, Eli Lilly, Glaxo Smith Kline, Lundbeck, Novartis, Servier, Phenicia, UPO. The Mexican National Comorbidity Survey (MNCS) is supported by The National Institute of Psychiatry Ramon de la Fuente (INPRFMDIES 4280) and by the National Council on Science and Technology (CONACyT-G30544- H), with supplemental support from the PanAmerican Health Organization (PAHO). The Northern Ireland Study of Mental Health was funded by the Health & Social Care Research & Development Division of the Public Health Agency. The Peruvian World Mental Health Study was funded by the National Institute of Health of the Ministry of Health of Peru. The Romania WMH study projects "Policies in Mental Health Area" and "National Study regarding Mental Health and Services Use" were carried out by National School of Public Health & Health Services Management (former National Institute for Research & Development in Health), with technical support of Metro Media Transilvania, the National Institute of Statistics-National Centre for Training in Statistics, SC. Cheyenne Services SRL, Statistics Netherlands and were funded by Ministry of Public Health (former Ministry of Health) with supplemental support of Eli Lilly Romania SRL. The South Africa Stress and Health Study (SASH) is supported by the US National Institute of Mental Health (R01-MH059575) and National Institute of Drug Abuse with supplemental funding from the South African Department of Health and the University of Michigan. DS is supported by the Medical Research Council of South Africa (MRC).

The Psychiatric Enquiry to General Population in Southeast Spain – Murcia (PEGASUS-Murcia) Project has been financed by the Regional Health Authorities of Murcia (Servicio Murciano de Salud and Consejería de Sanidad y Política Social) and Fundación para la Formación e Investigación Sanitarias (FFIS) of Murcia. The Ukraine Comorbid Mental Disorders during Periods of Social Disruption (CMDPSD) study is funded by the US National Institute of Mental Health (R01-MH61905). The US National Comorbidity Survey Replication (NCS-R) is supported by the National Institute of Mental Health (NIMH; U01-MH60220) with supplemental support from the National Institute of Drug Abuse (NIDA), the Substance Abuse and Mental Health Services Administration (SAMHSA), the Robert Wood Johnson Foundation (RWJF; Grant 044708), and the John W. Alden Trust.

None of the funders had any role in the design, analysis, interpretation of results, or preparation of this paper. The views and opinions expressed in this report are those of the authors and should not be construed to represent the views of the World Health Organization, other sponsoring organizations, agencies, or governments.

A complete list of all within-country and cross-national WMH publications can be found at <http://www.hcp.med.harvard.edu/wmh/>

### **Declaration of Interest**

In the past 3 years, RCK received support for his epidemiological studies from Sanofi Aventis; was a consultant for Johnson & Johnson Wellness and Prevention, Shire, Takeda; and served on an advisory board for the Johnson & Johnson Services Inc. Lake Nona Life Project. Kessler is a co-owner of DataStat, Inc., a market research firm that carries out healthcare research.

In the past 3 years, KD has received personal fees from Lundbeck, Servier and Johnson & Johnson.

In the past 3 years, DJS has received research grants and/or consultancy honoraria from AMBRF/The Foundation for Alcohol Research, Biocodex, Cipla, Lundbeck, National Responsible Gambling Foundation, Novartis, Servier, and Sun.

In the past 3 years, NK has received support (in the form of consultancy fees, lecture fees and royalties) from Aishin-Seiki, EAP Consulting, Igaku-Shoin, Japan Dental Association, Japan Housing Finance Agency, Japan Productivity Center, Junpukai Health Care Center, Meiji, Nanko-do, Nanzan-do, Occupational Health Foundation, Osaka Chamber of Commerce and Industry, Otsuka and Sekisui Chemicals, PHP Publication and Taishu-kan. He has received research grants from Infocom Ltd., Japan Management Association, Japanese Ministry of Education, Science, and Technology, Japanese Ministry of Health, Labor and Welfare and SoftBank Corp. Fujitsu Software Technologies Ltd., has provided support to NK in the form of grants and royalties.

### **Ethical Standards**

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

## References

1. Brewin CR, Andrews B, Valentine JD. Meta-analysis of risk factors for posttraumatic stress disorder in trauma-exposed adults. *Journal of Consulting and Clinical Psychology*. 2000; 68: 748-66.
2. Koenen KC, Moffitt TE, Poulin R, Martin J, Caspi A. Early childhood factors associated with the development of post-traumatic stress disorder: results from a longitudinal birth cohort. *Psychological Medicine*. 2007; 37: 181-92.
3. Scott KM, Smith DR, Ellis PM. Prospectively ascertained childhood maltreatment and its associations with DSM-IV mental disorders in young adults. *Archives of General Psychiatry*. 2010; 67: 712-9.
4. Bremner JD, Southwick SM, Johnson DR, Yehuda R, Charney DS. Childhood physical abuse and combat-related posttraumatic stress disorder in Vietnam veterans. *American Journal of Psychiatry*. 1993; 150: 235-9.
5. Schnurr PP, Lunney CA, Sengupta A. Risk factors for the development versus maintenance of posttraumatic stress disorder. *Journal of Traumatic Stress*. 2004; 17: 85-95.
6. McLaughlin KA, Conron KJ, Koenen KC, Gilman SE. Childhood adversity, adult stressful life events, and risk of past-year psychiatric disorder: A test of the stress sensitization hypothesis in a population-based sample of adults. *Psychological Medicine*. 2010; 40: 1647-58.
7. Myers B, McLaughlin KA, Wang S, Blanco C, Stein DJ. Associations between childhood adversity, adult stressful life events, and past-year drug use disorders in the National Epidemiological Study of Alcohol and Related Conditions (NESARC). *Psychology of Addictive Behaviors*. 2014; 28: 1117-26.
8. Roberts AL, McLaughlin KA, Conron KJ, Koenen KC. Adult stressors, history of childhood adversity, and risk of perpetration of intimate partner violence among men and women. *American Journal of Preventive Medicine*. 2010; 40: 128-38.
9. Keyes KM, Shmulewitz D, Greenstein E, McLaughlin KA, Wall M, Aharonovich E, et al. Exposure to the Lebanon War of 2006 and effects on alcohol use disorders: The moderating role of childhood maltreatment. *Drug and Alcohol Dependence*. 2014; 134: 296-303.
10. Hammen C, Henry R, Daley SE. Depression and sensitization to stressors among young women as a function of childhood adversity. *Journal of Consulting and Clinical Psychology*. 2000; 68: 782-7.
11. Harkness KL, Bruce AE, Lumley MN. The role of childhood abuse and neglect in the sensitization to stressful life events in adolescent depression. *Journal of Abnormal Psychology*. 2006; 115: 730-41.
12. Espejo EP, Hammen C, Connolly NP, Brennan PA, Najman JM, Bor W. Stress sensitization and adolescent depressive severity as a function of childhood adversity: A link to anxiety disorders. *Journal of Abnormal Child Psychology*. 2006; 35: 287-99.
13. Kessler RC, Davis CG, Kendler KS. Childhood adversity and adult psychiatric disorder in the US National Comorbidity Survey. *Psychological Medicine*. 1997; 27: 1101-19.
14. Kessler RC, McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky AM, et al. Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *British Journal of Psychiatry*. 2010; 197: 378-85.
15. Green JG, McLaughlin KA, Berglund P, Gruber MJ, Sampson NA, Zaslavsky AM, et al. Childhood adversities and adult psychopathology in the National Comorbidity Survey

- Replication (NCS-R) I: Associations with first onset of DSM-IV disorders. *Archives of General Psychiatry*. 2010; 62: 113-23.
16. McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky A, Kessler RC. Childhood adversities and first onset of psychiatric disorders in a national sample of adolescents. *Archives of General Psychiatry*. 2012; 69: 1151-60.
  17. Breslau N, Kessler RC, Chilcoat HD, Schultz LR, Davis GC, Andreski P. Trauma and posttraumatic stress disorder in the community: The 1996 Detroit Area Survey of Trauma. *Archives of General Psychiatry*. 1998; 55: 626-32.
  18. Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB. Posttraumatic stress disorder in the National Comorbidity Survey. *Archives of General Psychiatry*. 1995; 52: 1048-60.
  19. Liu H, Petukhova M, Sampson NA, Aguilar-Gaxiola S, Alonso J, Andrade LH, et al. Association of DSM-IV post-traumatic stress disorder with traumatic experience type and history in the World Health Organization World Mental Health Surveys. *JAMA Psychiatry*. 2017; 74: 270-81.
  20. The World Bank. Data: Countries and Economies. The World Bank Group, 2016.
  21. Heeringa SG, Wells EJ, Hubbard F, Mneimneh ZN, Chiu WT, Sampson NA, et al. Sample designs and sampling procedures. In: *The WHO World Mental Health Surveys: Global perspectives on the epidemiology of mental disorders* (eds RC Kessler, TB Üstun): 14-32. Cambridge University Press, 2008.
  22. Harkness J, Pennell B-P, Villar A, Gebler N, Aguilar-Gaxiola S, Bilgen I. Translation procedures and translation assessment in the World Mental Health Survey Initiative. In: *The WHO World Mental Health Surveys: Global perspectives on the epidemiology of mental disorders* (eds RC Kessler, TB Üstun): 91-113. Cambridge University Press, 2008.
  23. Benjet C, Bromet E, Karam EG, Kessler RC, McLaughlin KA, Ruscio AM, et al. The epidemiology of traumatic event exposure worldwide: results from the World Mental Health Survey Consortium. *Psychological Medicine*. 2016; 46: 327-43.
  24. Kessler RC, Üstun TB. The World Mental Health (WHM) Survey Initiative Version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *International Journal of Methods in Psychiatric Research*. 2004; 13: 93-121.
  25. McLaughlin KA, Koenen KC, Friedman MJ, Ruscio AM, Karam EG, Shahly V, et al. Subthreshold posttraumatic stress disorder in the World Health Organization World Mental Health Surveys. *Biological Psychiatry*. 2015; 77: 375-84.
  26. Haro JM, Arbabzadeh-Bouchez S, Brugha TS, di Girolamo G, Guyer M, Jin R, et al. Concordance of the Composite International Diagnostic Interview Version 3.0 (CIDI 3.0) with standardized clinical assessments in the WHO World Mental Health Surveys. *International Journal of Methods in Psychiatric Research*. 2006; 15: 167-80.
  27. Gardner M, Altman D. *Statistics with Confidence: Confidence Intervals and Statistical Guidelines*. BMJ Books, 2000.
  28. Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*. 1998; 14: 245-58.
  29. Straus MA. Measuring intrafamily conflict and violence: the Conflict Tactics (CT) Scales. *Journal of Marriage and the Family*. 1979; 41: 75-88.
  30. Courtney ME, Piliavin I, Grogan-Kaylor A, Nesmith A. Foster youth transitions to adulthood: A longitudinal view of youth leaving care. Institute for Research on Poverty, 1998.

31. Endicott J, Andreasen N, Spitzer RL. Family History Research Diagnostic Criteria. Biometrics Research, NY State Psychiatric Institute, 1978.
32. Kendler KS, Silberg JL, Neale MC, Kessler RC, Heath AC, Eaves LJ. The family history method: whose psychiatric history is measured? *American Journal of Psychiatry*. 1991; 148: 1501-4.
33. Klebanoff MA, Cole SR. Use of multiple imputation in the epidemiologic literature. *American Journal of Epidemiology*. 2008; 168: 355-7.
34. Wolter KM. Introduction to variance estimation. Springer-Verlag, 1985.
35. SAS Institute Inc. SAS Software Version 9.2. 2008.
36. McLaughlin KA, Gademann AM, Hwang I, Sampson NA, Al-Hamzawi A, Andrade LH, et al. Parent psychopathology and offspring mental disorders: results from the WHO World Mental Health Surveys. *Br J Psychiatry*. 2012; 200(4): 290-9.
37. Caspi A, Houts RM, Belsky DW, Goldman-Mellor SJ, Harrington H, Israel S, et al. The p factor: One general psychopathology factor in the structure of psychiatric disorders? *Clinical Psychological Science*. 2014; 2: 119-37.
38. Rudolph KD, Flynn M. Childhood adversity and youth depression: Influence of gender and pubertal status. *Development and Psychopathology*. 2007; 19(497-521).
39. Breslau N, Koenen KC, Luo Z, Agnew-Blais J, Swanson S, Houts RM, et al. Childhood maltreatment, juvenile disorders and adult post-traumatic stress disorder: a prospective investigation. *Psychological Medicine*. 2014; 44: 1937-45.
40. Moore SA, Zoellner LA. Overgeneral autobiographical memory and traumatic events: an evaluative review. *Psychological Bulletin*. 2007; 133: 419-37.
41. Heleniak C, Jenness J, Van der Stoep A, McCauley E, McLaughlin KA. Childhood maltreatment exposure and disruptions in emotion regulation: A transdiagnostic pathway to adolescent internalizing and externalizing psychopathology. *Cognitive Therapy and Research*. 2016; 40: 394-415.
42. Hennessy KD, Rabideau GJ, Cicchetti D, Cummings EM. Responses of physically abused and nonabused children to different forms of interadult anger. *Child Development*. 1994; 65: 815-28.
43. Wichers M, Schrijvers D, Geschwind N, Jacobs N, Myin-Germeys I, Thiery E, et al. Mechanisms of gene-environment interactions in depression: evidence that genes potentiate multiple sources of adversity. *Psychological Medicine*. 2009; 39: 1077-86.
44. McCrory EJ, De Brito SA, Kelly PA, Bird G, Sebastian CL, Mechelli A, et al. Amygdala activation in maltreated children during pre-attentive emotional processing. *British Journal of Psychiatry*. 2013; 202: 269-76.
45. McCrory EJ, De Brito SA, Sebastian CL, Mechelli A, Bird G, Kelly PA, et al. Heightened neural reactivity to threat in child victims of family violence. *Current Biology*. 2011; 21: R947-8.
46. Suzuki H, Luby JL, Botteron KN, Dietrich R, McAvoy MP, Barch DM. Early life stress and trauma and enhanced limbic activation to emotionally valenced faces in depressed and healthy children. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2014; 53: 800-13.
47. McLaughlin KA, Peverill M, Gold AL, Alves S, Sheridan MA. Child maltreatment and neural systems underlying emotion regulation. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2015; 54: 753-62.

48. Feldman R, Granat A, Pariente C, Kanety H, Kuint J, Gilboa-Schechtman E. Maternal depression and anxiety across the postpartum year and infant social engagement, fear regulation, and stress reactivity. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2009; 48: 919-27.
49. Davis EP, Glynn LM, Schetter CD, Hobel C, Chicz-Demet A, Sandman CA. Prenatal exposure to maternal depression and cortisol influences infant temperament. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2007; 46: 737-46.
50. McLaughlin KA, Busso DS, Duys A, Green JG, Alves S, Way M, et al. Amygdala response to negative stimuli predicts PTSD symptom onset following a terrorist attack. *Depression and Anxiety*. 2014; 10: 834-42.
51. Admon R, Lubin G, Stern O, Rosenberg K, Sela L, Ben-Ami H, et al. Human vulnerability to stress depends on amygdala's predisposition and hippocampal plasticity. *Proceedings of the National Academy of Sciences*. 2009; 106: 14120-5.
52. Mather M, Canli T, English T, Whitfield S, Wais P, Ochsner K, et al. Amygdala responses to emotionally valenced stimuli in older and younger adults. *Psychological Science*. 2004; 15: 259-63.
53. McLaughlin KA, Sheridan MA, Lambert HK. Childhood Adversity and Neural Development: Deprivation and Threat as Distinct Dimensions of Early Experience. *Neuroscience and Biobehavioral Reviews*. 2014; 47: 578-91.
54. Jenness J, Jager-Hyman S, Heleniak C, Beck AT, Sheridan MA, McLaughlin KA. Catastrophizing, rumination, and reappraisal prospectively predict adolescent PTSD symptom onset following a terrorist attack. *Depression and Anxiety*. in press.
55. Kuyken W, Watkins E, Holden E, Cook W. Rumination in adolescents at risk for depression. *Journal of Affective Disorders*. 2006; 96: 39-47.
56. Sheikh MA, Abelsen B, Olsen JA. Clarifying associations between childhood adversity, social support, behavioral factors, and mental health, health, and well-being in adulthood: A population-based study. *Frontiers in Psychology*. 2016; 7.
57. Hardt J, Rutter M. Validity of adult retrospective reports of adverse childhood experiences: review of the evidence. *Journal of Child Psychology and Psychiatry*. 2004; 45: 260-73.
58. Widom CS, Shepard RL. Accuracy of adult recollections of childhood victimization: Part 1. Childhood physical abuse. *Psychological Assessment*. 1996; 8: 412-21.
59. Widom CS, Morris S. Accuracy of adult recollections of childhood victimization: Part 2. Childhood sexual abuse. *Psychological Assessment*. 1997; 9: 34-46.



**Table 1. Prevalence of childhood adversities (CAs) in the WHO World Mental Health Surveys (n=27,017)**

	Proportion of TEs occurring to respondents with a history of each CA <sup>1</sup>		Proportion of respondents with a history of each CA <sup>2</sup>	
	%	(SE)	%	(SE)
Maladaptive Family Functioning (MFF) CAs				
Parent Psychopathology	13.7	(0.6)	9.5	(0.3)
Parent Substance Abuse	7.8	(0.4)	6.0	(0.2)
Parent Criminality	6.5	(0.4)	4.0	(0.2)
Family Violence	16.3	(0.6)	9.7	(0.3)
Physical Abuse	20.9	(0.7)	10.7	(0.3)
Sexual Abuse	3.8	(0.2)	1.9	(0.1)
Neglect	8.2	(0.3)	5.7	(0.2)
Other CAs				
Parent Death	14.4	(0.5)	13.3	(0.3)
Parent Divorce	9.7	(0.5)	8.0	(0.3)
Other Parent Loss	7.7	(0.4)	6.0	(0.2)
Serious Physical Illness	6.3	(0.4)	3.5	(0.2)
Economic Adversity	6.1	(0.3)	4.7	(0.2)

MFF, maladaptive family functioning; CAs, childhood adversities; TE, traumatic event

<sup>1</sup> Given that randomly-selected were weighting by the inverse of their probabilities of selection and then multiplied by the Part II weight, the weighted sample of traumatic events (TEs) represents the population of all TEs experienced in the population.

<sup>2</sup> Prevalence of CAs among the 27,017 respondents included in the analysis.

**Table 2. Multivariate associations (odds-ratios) between CAs and PTSD in response to a randomly-selected traumatic event (TE) in the WHO World Mental Health Surveys (n=27,017)<sup>1</sup>**

	M1 <sup>2</sup>		M2 <sup>3</sup>		M3 <sup>4</sup>		M4 <sup>5</sup>		M5 <sup>6</sup>		M6 <sup>7</sup>		M7 <sup>8</sup>	
	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
Maladaptive Family Functioning (MFF) CAs														
Parent Psychopathology	2.2*	(1.6-3.0)	2.0*	(1.5-2.7)	--	--	--	--	--	--	--	--	--	--
Parent Substance Abuse	1.2	(0.9-1.7)	0.9	(0.6-1.3)	--	--	0.9	(0.6-1.3)	0.9	(0.6-1.3)	0.9	(0.6-1.3)	--	--
Parent Criminality	1.4	(0.9-2.3)	1.1	(0.7-1.9)	--	--	1.1	(0.7-1.9)	1.1	(0.7-1.9)	1.1	(0.7-1.9)	--	--
Family Violence	1.1	(0.7-1.7)	1.0	(0.6-1.5)	--	--	1.0	(0.6-1.5)	1.1	(0.7-1.7)	1.1	(0.7-1.7)	--	--
Physical Abuse	1.6*	(1.1-2.3)	1.3	(0.9-2.0)	--	--	0.7	(0.4-1.1)	--	--	--	--	--	--
Sexual Abuse	2.0*	(1.2-3.4)	1.8*	(1.1-3.0)	--	--	0.9	(0.5-1.5)	--	--	--	--	--	--
Neglect	2.2*	(1.4-3.2)	1.9*	(1.2-2.8)	--	--	0.9	(0.6-1.5)	--	--	--	--	--	--
$\chi^2_{7}$	--		40.8*				3.4		$\chi^2_{3} = 0.3$		$\chi^2_{3} = 0.3$			
$\chi^2_{6}$			18.5*											
Other CAs														
Parent Death	1.4	(0.9-2.4)	1.4	(0.8-2.3)	--	--	1.4	(0.8-2.4)	1.4	(0.7-2.4)	1.4	(0.7-2.4)	--	--
Parent Divorce	0.9	(0.6-1.2)	0.9	(0.6-1.2)	--	--	0.9	(0.6-1.3)	0.8	(0.6-1.3)	0.9	(0.6-1.3)	--	--
Other Parent Loss	0.9	(0.6-1.4)	0.9	(0.5-1.2)	--	--	0.7	(0.4-1.3)	0.7	(0.4-1.3)	0.7	(0.4-1.3)	--	--
Serious Physical Illness	1.1	(0.8-1.7)	1.1	(0.7-1.5)	--	--	1.0	(0.7-1.6)	1.0	(0.6-1.6)	1.0	(0.7-1.6)	--	--
Economic Adversity	1.3	(0.8-2.0)	1.3	(0.7-1.8)	--	--	1.1	(0.6-1.9)	1.1	(0.6-2.0)	1.1	(0.6-2.0)	--	--
$\chi^2_{5}$	--		5.8		--		6.1		6.7		6.8			
$\chi^2_{4}$			5.8		--		5.8		6.3		6.4			
$\chi^2_{12}$	--		63.8*		--		9.3		6.9		6.9			
Number of MFF CAs														
0-4 <sup>9</sup>	--	--	--	--	1.7*	(1.5-2.1)	2.0*	(1.5-2.7)	1.8*	(1.5-2.1)	--	--	1.8*	(1.5-2.1)
1	--	--	--	--	--	--	--	--	--	--	1.6*	(1.1-2.2)	--	--
2	--	--	--	--	--	--	--	--	--	--	2.8*	(1.8-4.4)	--	--
3	--	--	--	--	--	--	--	--	--	--	5.4*	(2.7-10.6)	--	--
4	--	--	--	--	--	--	--	--	--	--	13.7*	(3.6-51.4)	--	--
$\chi^2_{4}$	--		--		--		--		--		36.3*			
$\chi^2_{3}$	--		--		--		--		--		19.6*			
Number of Other CAs														
1	--	--	--	--	1.1	(0.8-1.6)	--	--	--	--	--	--	--	--
2+	--	--	--	--	1.1	(0.8-1.7)	1.1	(0.6-2.1)	1.1	(0.6-2.2)	1.1	(0.6-2.2)	--	--
$\chi^2_{2}$	--		--		0.4		--		--		--			
Mean-squared error <sup>10</sup>			0.039214		0.039162		0.039232		0.039206		0.039278		0.039146	
AUC <sup>10</sup>			0.738		0.741		0.737		0.737		0.734		0.743	

MFF, maladaptive family functioning; CAs, childhood adversities; TE, traumatic event

\*Significant at the .05 level, two-sided test

<sup>1</sup>Models were estimated using logistic regression with PTSD in response to a randomly-selected TE as the outcome variable. Each model controlled for survey, age, sex, 9 dummy variables for TE type, and 6 dummy variables for prior exposure to the same TE type for trauma types shown in prior analysis to influence later risk of PTSD in response to a TE of the same type (see citation 19 for details)

<sup>2</sup>Models were estimated with one CA at a time and the controls noted in footnote 1.

<sup>3</sup>The model was estimated with all 12 CAs and the controls noted in footnote 1.

<sup>4</sup>The model was estimated with dummy variables for the number of CAs without any information about CA type and the controls in footnote 1.

<sup>5</sup>The model was estimated with dummy variables for both number and type of CAs plus the controls noted in footnote 1.

<sup>6</sup>The model was estimated with dummy variables for both number and type of CAs, removing the variables for CA type for the 4 MFF CAs included in the 0-4 count variable (see footnote 9), plus the controls noted in footnote 1.

<sup>7</sup>The model was estimated with dummy variables for both number and type of CAs, removing the variables for CA type for the 4 MFF CAs included in the 0-4 count variable (see footnote 9), plus the controls noted in footnote 1.

<sup>8</sup>The model was estimated with the 0-4 count variable (see footnote 9), plus the controls noted in footnote 1.

<sup>9</sup>The 0-4 variable is a count of the number of the following CAs experienced by the respondent: physical abuse, sexual abuse, neglect, and parent psychopathology.

<sup>10</sup>These measures are based on replicated 10-fold cross-validation with 10 replications.

**Table 3. Variation in the associations (odds-ratios) of CAs and PTSD in response to a randomly-selected traumatic event (TE) by TE type and age of TE exposure in the WHO World Mental Health Surveys (n=27,017)<sup>1</sup>**

	Variation by TE Type <sup>2</sup>		Variation by Age of TE Exposure <sup>3</sup>	
	OR	(95% CI)	OR	(95% CI)
Interaction of Number of MFF CAs with TE Types				
Civilian in a war zone	0.8	(0.3-2.3)	--	--
Civilian in a region of terror	1.1	(0.7-2.0)	--	--
Kidnapped	3.1*	(1.2-7.7)	--	--
Witnessed death	1.6*	(1.0-2.6)	--	--
Saw atrocities	0.4	(0.1-1.6)	--	--
Sexual violence	1.2	(0.8-1.8)	--	--
Natural disaster	0.3*	(0.1-0.9)	--	--
Unexpected death of a loved one	1.4	(0.9-2.3)	--	--
All other TEs	1.2	(0.8-1.7)	--	--
$\chi^2_8$		13.9		--
Interaction of Number of MFF CAs with Age of TE Exposure				
0-12 years	--	--	1.6*	(1.1-2.3)
13-24 years	--	--	1.8*	(1.3-2.6)
25-44 years	--	--	1.9*	(1.3-2.7)
45-59 years	--	--	1.5	(0.9-2.6)
60+	--	--	0.1*	(0.0-0.3)
$\chi^2_4$		--		16.5*
Interaction of Number of MFF CAs with Country-Income Group				
High	--	--	--	--
Other	--	--	--	--
$\chi^2_1$		--		--

MFF, maladaptive family functioning; CAs, childhood adversities; TE, traumatic event

\*Significant at the .05 level, two-sided test

<sup>1</sup>See Footnote 1 in Table 2 for a description of the overall modeling approach. CA and control variables are based on the best-fitting model of CAs in predicting PTSD in response to a randomly-selected TE (M7, Table 1).

<sup>2</sup>The model included 9 dummy variables for the interaction of the 0-4 MFF CA count variable with TE types in addition to CA and control variables from M7, Table 2. TE types are based on prior factor analysis in this sample (see Methods section for details).

<sup>3</sup>The model included 5 dummy variables for the interaction of the 0-4 MFF CA count variable with age at TE exposure in addition to CA and control variables from M7, Table 2.

<sup>4</sup>The model included 2 dummy variables for the interaction of the 0-4 MFF CA count variable with country-income group in addition to CA and control variables from M7, Table 2.