

Changes in physical activity and sedentary behaviours from before to during the COVID-19 pandemic lockdown: A Systematic Review

Stephanie Stockwell,^{1†} Mike Trott,^{1,2†} Mark Tully,³ Jae Shin,⁴ Yvonne Barnett,⁵ Laurie Butler,⁶ Daragh McDermott,⁶ Felipe Schuch,⁷ Lee Smith¹

¹Cambridge Centre for Sport and Exercise Sciences, Anglia Ruskin University - Cambridge Campus, Cambridge, UK

²Vision and Eye Research Institute, Anglia Ruskin University, Cambridge, UK

³Institute of Mental Health Sciences, School of Health Sciences, University of Ulster, Newtownabbey, UK

⁴Department of Pediatrics, Yonsei University College of Medicine, Seodaemun-gu, Seoul, Republic of Korea

⁵Anglia Ruskin University - Cambridge Campus, Cambridge, UK ⁶ Faculty of Science and Engineering, Anglia Ruskin University, Cambridge, UK

⁷Department of Sports Methods and Techniques, Federal University of Santa Maria, Santa Maria, Brazil

*Corresponding Author

Stephanie Stockwell, The Cambridge Centre for Sport and Exercise Sciences, Anglia Ruskin University, Cambridge UK, CB1 1PT; Stephanie.Stockwell1@pgr.anglia.ac.uk

†Contributed equally to the work

Abstract

Objective

In March 2020, several countries banned unnecessary outdoor activities during COVID-19, commonly called 'lockdowns. These lockdowns have the potential to impact associated levels of physical activity and sedentary behaviour. Given the numerous health outcomes associated with physical activity and sedentary behaviour, the aim of this review was to summarise literature that investigated differences in physical activity and sedentary behaviour before vs during the COVID-19 lockdown.

Design, data sources and eligibility criteria

Electronic databases were searched from November 2019 to October 2020 using terms and synonyms relating to physical activity, sedentary behaviour and COVID-19. The coprimary outcomes were changes in physical activity and/or sedentary behaviour captured via device-based measures or self-report tools. Risk of bias was measured using the Newcastle-Ottawa Scale.

Results

Sixty six articles met the inclusion criteria and were included in the review (total n=86 981). Changes in physical activity were reported in 64 studies, with the majority of studies reporting decreases in physical activity and increases in sedentary behaviours during their respective lockdowns across several populations, including children and patients with a variety of medical conditions.

Conclusion

Given the numerous physical and mental benefits of increased physical activity and decreased sedentary behaviour, public health strategies should include the creation and implementation of interventions that promote safe physical activity and reduce sedentary behaviour should other lockdowns occur.

Introduction

In March 2020, WHO declared the COVID-19 outbreak a global pandemic, and as of 26 October 2020, over 42 000 000 confirmed cases have been diagnosed in more than 130 countries and territories, resulting in approximately 1 150 000 deaths.¹ COVID-19 has led to over 100 countries enforcing social distancing to reduce the rate of COVID-19 transmission, commonly called 'lockdown'.² The severity of lockdown has varied from country to country, even region to region, with some countries limiting the distance people could travel from their homes, and some banning any unnecessary outdoor activity.² These lockdowns have impacted people's work, education, travel and recreation, and subsequent levels of physical activity (PA) and sedentary behaviours (SB).³

PA can be defined as any bodily movement produced by skeletal muscle that results in energy expenditure,⁴ and can include exercising, walking, gardening and doing household chores. Research shows that PA is positively associated with several desirable outcomes, including social contentedness,⁵ physical health⁶ and mental health.^{7,8} Specific to COVID-19, PA has been shown to improve physical and mental health and has been suggested to provide protective elements against COVID-19.⁹⁻¹¹ Furthermore, it has been reported that the COVID-19 lockdown yielded decreases in PA,¹² however, the literature has not been systematically reviewed to date.

SB can be defined as any waking behaviour with an energy expenditure of ≤ 1.5 Metabolic Equivalents (METs) while in a sitting or reclining posture,¹³ including watching TV, video gaming and computer use. The literature has shown SB to be negatively associated with physical, mental health and social outcomes.^{14,15} Specific to COVID-19, it has been reported that periods of

enforced quarantine can yield increases in SB,^{9-13,16} however, this has not been systematically assessed to date in the context of the COVID-19 lockdown.

Understanding the changes in PA and SB behaviours during lockdown is important not only for health outcomes associated with these behaviours, but also for aiding development of public health interventions in specific populations (such as PA promotion and interventions to decrease SB) should another lockdown be enforced, a similar pandemic scenario and/or during the return to 'normal life'.¹⁷ The aim of this study, therefore, was to conduct a comprehensive systematic review on changes in all reported PA and SB behaviours during versus before the COVID-19 pandemic lockdown, stratifying between adults and children, and special populations.

Methods

The current systematic review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.¹⁸ Details of the full protocol for this systematic review were registered on PROSPERO (protocol number: CRD42020193065).

Search strategy

Electronic databases were searched from November 2019 to June 2020 including PubMed, EMBASE, PsycINFO, CINAHL, Social Science Citation Index, Cochrane Central Register of Controlled Trials, SPORTDiscus and Scopus. Grey literature was searched by entering terms into OpenGrey. Search terms were as follows:

COVID-19 OR "Novel Coronavirus" OR "2019 novel coronavirus" OR 2019-nCoV OR SARS-CoV-2

AND isolation OR lock* OR self-isolation

AND

“Physical activity” OR exercise OR walking OR running OR cycling OR swimming OR sports OR sedentary OR “sedentary behaviour” OR activity OR “screen time” OR sitting.

Full information on database-specific search strategies can be found in online supplemental table 1.

Results of the searches were included in a bibliographic database and duplicates removed. Titles and abstracts of the retrieved studies were screened for inclusion by two reviewers independently (SS and MT), and then the full text of all potentially eligible papers was reviewed independently by the same reviewers before making a final decision on eligibility. Any discrepancies were discussed until a decision was achieved. A third senior reviewer acted as an adjudicator if a decision was not reached (LS).

Study inclusion and exclusion

Studies were included if they met the following criteria: (1) observational cross-sectional, prospective or retrospective cohort studies (2) that investigate any form of PA and/or SB (as defined by the authors) (3) in any population (healthy or with a specific disease condition) (4) before and during the COVID-19 Lockdown (5) in any setting. Published articles that had received ethical approval from an ethics committee and were written in English were included. Studies

were excluded if they were not observational in design (eg, qualitative, primary randomised controlled trials, primary case series, editorials or commentaries or study protocols). Furthermore, the rapid publication of studies related to COVID-19 meant many bypassed the typical institutional ethical approval process; therefore, studies were excluded if they failed to explicitly include an explicit statement stating that institutional ethical approval was received. If no ethical approval was in the manuscript, corresponding authors were contacted to establish if institutional approval had been granted. If no reply was received, or institutional ethics was confirmed as being not obtained, these studies were excluded (see online supplemental table 2).

Primary outcomes

The coprimary outcomes were changes in PA and/or SB captured via device-based measures or self-report tools.

Data extraction

Data were extracted by two reviewers (SS and MT) independently including: first author, year, country, aims of the study, type of the study (pretest and post-test, cross-sectional, cohort), descriptions of the lockdown by the authors for the respective location of data collection, number of participants, participant characteristics (eg, age, sex), inclusion criteria, type of recruitment, type and definition of PA and SB investigated, type of measurement of PA and SB, confounding variables, acknowledged limitations by authors and authors conclusions, other/notes. A third reviewer (LS) was available to resolve any discrepancies. The data was synthesised in a narrative approach.

Quality assessment

Risk of bias was assessed by two independent reviewers (SS and MT) using the Newcastle-Ottawa Scale (NOS),¹⁹ later adapted for cross-sectional studies.²⁰ A third reviewer (LS) was available to resolve any inconsistencies. There are three parts in which studies are assessed and stars awarded: (1) selection (max. 5 stars)—representativeness of the sample, sample size, non-respondents and ascertainment of the exposure (risk factor); (2) comparability (max. 2 stars)—participants in different outcome groups are comparable; (3) outcome (max. 3 stars)—assessment of outcome, and statistical test. Scores can range from 0 to 10 stars, with higher scores indicating better quality research.

Results

After initial screening, 187 studies were eligible for full-text review. From these, 66 studies^{12 21–84} were eligible for inclusion. The PRISMA flow chart is shown in online supplemental figure 1. Full study characteristics can be found in online supplemental table 3. The 66 included studies yielded a total of 86 981 participants and the age ranged from 13 to 86 years old. Regarding specific populations, forty-five studies were conducted on healthy adults (four in specifically elite athletes and five in university students), and six studies in healthy children. Regarding populations with medical conditions, two studies were conducted on adult women with eating disorders, two respective studies with adult participants with type 1 and types 2 diabetes, one respective study with adult participants with ‘chronic medical conditions’, one study on heart failure patients, one study on neuromuscular disease, one study on obesity clinic patients, one study on participants with a ‘perceived risk’ of severe COVID-19 symptoms, one study on pregnant women and one study reported on children with obesity. The mean NOS score of the included studies was 4.8 (SD=1.0; range 3–7;). For detailed NOS scoring, see online supplemental table 4.




Figure 1: PRISMA Flowchart of included studies

PA in healthy adults

Forty-five studies examined PA changes in healthy adults, with only four studies^{31 75 80 85} using device-based measures of PA. The remaining 41 studies used subjective questionnaires, and in 30 studies these questionnaires were not previously validated. The majority of studies (26/45) reported PA changes in the form of time (eg, METS/min/week, mins/day or steps/day), with the

remaining studies reporting PA changes as a percentage of the respective population (see online supplemental table 5).

Of the studies that measured PA change in the form of time spent on PA, all but one study⁷⁰ reported overall decreases in the amount of PA pre-COVID-19 versus post-COVID-19 lockdown. When stratifying across different forms of PA, two studies^{27 28} reported increases in time spent in 'leisure-time PA' and one study⁸⁵ reported increases in time spent in 'endurance training' in elite cyclists, although total PA still decreased in all three studies. All other studies reported time spent in all subtypes of PA—for example, light, moderate, vigorous and walking—if specified) decreased. Of the studies that measured PA changes as a percentage of the respective populations, eight studies^{29 33 38 53 55 59 67 81} reported that >50% of the examined population decreased PA during lockdown, with all other studies reporting >50% of the examined population's PA either stayed the same or decreased. For further information, see online supplemental table 5.

PA in healthy children and adolescents

Of the six studies that examined PA changes in healthy children and adolescents, all were measured using subjective questionnaires, with half using validated questionnaires. Two studies^{48 84} used total scores from validated questionnaires and two studies^{65 82} reported PA changes in the form of a time measurement, all reporting decreases in PA. Two studies^{62 71} reported PA changes as a percentage of the respective population and reported >50% of the population decreased their PA during lockdown.

PA in adults and children with medical conditions

Thirteen studies examined populations with medical conditions for which all but one study,⁸⁰ used subjective measurements of PA change, and in only 6/12 were these previously validated measurement tools. Regarding the types of changes reported, nine studies^{23 26 36 40 43 44 72 80 86} reported changes in time spent in PA, all reporting decreases in PA time. The remaining four studies^{34 66 69 74} reported PA changes as a percentage of respective populations, with all reporting >50% of the population decreasing their PA during lockdown.

SB in healthy adults

Of the 26 studies examined changes in SBs, 18 were conducted in healthy adults. All studies used subjective questionnaires and validated questionnaires were used in six. Studies reported changes in SB as either time spent on SB or as a percentage of the sample. The majority of studies (13/18) reported SB changes in the form of time spent, with the remaining studies reporting SB changes as a percentage of the respective population. Increased SB was reported in all 26 studies. For further information, see online supplemental table 6.

SB in healthy children and adolescents

Of the five studies that measured changes in SB in children and adolescents, three studies^{42 62 65} used non-validated questionnaires and the remaining two studies^{71 82} used validated questionnaires. Time spent in SB was reported in 3/5 studies, with the remaining two studies reporting changes in SB as a percentage of their respective populations. All five studies reported increases in SB.

SB in adults and children with medical conditions

All of the three^{26 43 86} studies that measured changes in SB in special populations used non-validated questionnaires, and reported that time spent in SB increased during the lockdown.

Discussion

The current systematic review of 66 studies demonstrated that the majority of studies found that PA declined and SB increased during the COVID-19 pandemic lockdown, regardless of the subpopulation or the methodology used. In healthy adults and children, PA during lockdown decreased compared with prelockdown, despite various government organisations and health or exercise practitioners providing guidance on how to stay active during the pandemic and in self-quarantine.^{87–89} When stratifying between prelockdown PA levels, three studies found that people who were more active prelockdown were more likely to show larger decreases in PA.^{37 49} ⁶⁷ PA has also been consistently linked with several mental health conditions, suggesting that decreases in PA may lead to increases in undesirable mental health outcomes. Indeed, studies have shown significant increases in anxiety and depression levels during the lockdown.¹⁰ Given that decreases in PA have been shown to yield negative affect, increases in anxiety and lower energy levels,³⁹ PA promotion during lockdowns should be aimed not just at people who are currently sedentary, but also for those with high PA levels outside of lockdown. Due to the likelihood of further COVID-19-related restrictions (or another similar pandemic), the promotion of digital based PA (such as PA apps, online video fitness classes or physical training) is recommended. Digital based PA yielded favourable results during the first COVID-19 lockdown, with studies showing positive associations with such digital based initiatives and overall PA during a lockdown.⁸³

Another finding of this review was that participants who had medical conditions also yielded decreases in PA levels, except for patients with an eating disorder. The decreases in PA is particularly concerning as in several of the medical conditions studied because PA can be a form

of treatment or symptom alleviation. For example, levels of PA have been shown to be positively associated with quality of life outcomes in both type 1 and type 2 diabetes.^{90 91} Concurrently, increases in SB have been shown to yield detrimental outcomes in patients with these conditions, except for patients with eating disorder.^{92 93} Given these added risks of decreasing PA and increasing SB in these special populations, PA promotion and strategies to reduce SB should be implemented should further lockdowns occur. Moreover, practitioners working with these groups should be especially mindful of the detriment that decreasing PA and increasing SB could yield during lockdowns and make the monitoring of PA levels a priority. Patients with eating disorders were found to increase their PA, specifically exercise, during lockdowns. This is equally concerning as there is often pathological relationship between eating disorders and exercise and can lead to increased risks of physical complications such as stress fractures.⁹⁴ Therefore, practitioners working with patients with eating disorders are advised to keep closely monitoring patients as much as possible during future lockdowns.

There were also large decreases in both the training volume and training intensity of elite athletes while in lockdown, which has led to relative decreases in sport-specific physical performance tests post-lockdown.⁸⁵ This decrease in athletic readiness for competition should be noted and considered by practitioners who are working with elite athletes, especially regarding training loads and competition scheduling postlockdown.

According to the behavioural change wheel, for a behaviour—for example, PA or SB—to occur, there are three components that are required: capability (psychological and physical), opportunity (physical and social) and motivation (reflective and automatic).⁹⁵ Despite information on safe exercise during lockdown being available from exercise professionals and some governments

(psychological capability), it is not clear from the included studies the reasons why people did or did not engage in PA; however, we can speculate potential reasons for these findings. A reduction in PA is expected as lockdowns required that governments closed sport and leisure facilities, group activities were suspended, and in many countries limits were in place for time spent outdoors.⁹⁶ This potentially meant people's regular PA routines were difficult to continue with during lockdown, as indicated by the evidence stating that people considerably changed their modes of PA during lockdown.⁹⁷ For example, one study found that all types of PA decreased except for 'moderate intensity leisure-time PA' (such as housework and gardening) increased,²⁶ another found that 'yard work' increased,⁷⁷ and another found that 'housework' increased during lockdown.⁸² However, despite these mode-specific increases, total PA levels in these respective populations still decreased. This suggests that promoting increases in house-related PA may not be sufficient to increase total PA during lockdowns.

There was also an increase in the number of people working from home during lockdown,⁹⁸ consequently, PA ordinarily accumulated during commuting will have substantially decreased. A previous study found that adults in the UK (mean age 50.5 years) accumulated 195 min/week (± 188.6) of active travel.⁹⁹ Those who actively commute report significantly greater total PA than those who do not, despite no significant differences in recreational PA shown.^{99 100} In addition, with schools closed, many parents were balancing home schooling, while working from home themselves; in the UK, this was the case for 85% of employees with school-aged children.¹⁰¹ A decrease in opportunities to be active and additional responsibilities may have led to a decrease in PA.

The majority of the studies in this review showed increases in SB during lockdown. This is unsurprising as many people worked from home, leading to extended sedentary periods and increased screen time.^{102 103} For instance, de Haas et al¹⁰⁴ reported that 44% of Dutch workers had either started to work from home or increased their home working hours, with 30% reporting increases in remote meetings (eg, via videoconferencing). In addition, with most gyms, leisure and sporting facilities closed, time allowed outdoors limited or not allowed, some people may have found it difficult to be active during the lockdown.^{3 105} With increased 'free' time, many may resort to engaging in pastimes such as reading, playing video games and watching television (TV), many of which are sedentary.⁶⁷

Given that the majority of studies reported a decrease in PA with a concurrent increase in SB during the lockdown, and the impact of these on physical and mental health, it is recommended that interventions or policies are implemented to increase PA (eg, body weight home-workouts, using online exercise classes, walking, running and cycling outdoors) and decrease SB (eg, by using a standing desk and taking regular breaks from sitting) should further lockdowns be enforced in the future. In addition, interventions for PA and/or SB postlockdown should consider that individuals may suffer deconditioning as a result of the lockdowns.

Many of the included studies used surveys to gather information about 'exercise', 'PA', 'sport' and 'training' but failed to report on how these terms were defined to participants. Future studies should report these definitions for clarity and comparison to be made more easily between studies. This lack of definition may mean that despite 'exercise' and 'training' decreasing, changes in daily PA may be different in these studies. Monbiot¹⁰⁶ reports volunteers providing food packages, collecting medical supplies for the elderly, providing childcare for those in need,

meaning they potentially accumulate similar or more 'activity' than they realise as it is not prescribed 'exercise' or 'training'.

It is important to note different degrees of lockdown in different countries, even regions within a country, across different dates occurred, making it difficult to quantify the severity of a lockdown and therefore challenging to objectively assess how this impacted behaviours. For instance, those in countries that were able to exercise outdoors following social distancing guidelines may have engaged differently in PA/SB behaviours to those who were not able to leave home, despite both countries being in 'lockdown'. Although the authors have presented the lockdown descriptions for each included study as reported by the authors, these descriptions vary greatly in detail, making it challenging to categorise them into 'levels' of lockdown. The creation of a scale to indicate lockdown severity would be highly beneficial for comparisons to be made between countries when investigating different behaviours, or at the very least it is recommended that this type of information is reported in all future studies. Moreover, within countries some people are given specific guidance (eg, shielding) which requires more intensive lockdown than the general population—none of the included studies recorded this information. It may be beneficial to know participants' adherence to lockdown guidelines to provide an indication of potential opportunity to engage in PA. Most studies also report PA without investigating in detail the types, intensities and durations of PA engaged in before and during lockdown, thus, it would be beneficial to investigate these as the magnitude of changes will impact the effects on health.

Limitations

While this systematic review is the first to our knowledge to assess changes in the frequency and modes of PA and SB prelockdown versus during the COVID-19 lockdown, the findings should be

considered within the limitations of the study. First, the tools used to measure PA and SB were highly heterogeneous, making direct comparison of respective results difficult. Second, demographic information was largely limited, meaning that we were unable to assess any further changes according to demographics further than the discussed topics, which would have given more insight into the review. In addition, the vast majority of studies were based on subjective questionnaires, which carry with them inherent limitations.¹⁰⁷ Moreover, many studies asked participants retrospectively about their prelockdown behaviours and their current behaviours during lockdown, thus, the accuracy of participants abilities to accurately recall their behaviours may be questionable. Lastly, most of the studies included were not designed to be nationally representative, making the generalisation of these results difficult.

Future research in this area should focus on yielding directly comparable data using validated PA and SB questionnaires or using objective accelerometer data where possible. In addition, it would be beneficial to have more detailed demographic information, information on the severity of lockdown and participant adherence to lockdown guidelines, and more detailed information on PA behaviours, for instance, the types, intensities and duration of PA before and during lockdown. Future research should also consider investigating the magnitude of the decrease in PA and increases in SB across different populations during the lockdown to aid the identification of populations most in need of targeted interventions. Lastly, future research should consider investigating the reasons why people are showing changes in PA and/or SB. Using behavioural change theory to assess barriers and facilitators to PA/SB during lockdowns would be highly beneficial in the creation of future interventions and policies should lockdowns occur in the future.

Conclusion

During the COVID-19 lockdown, PA levels have significantly reduced with concurrent increases in SB. Considering the evidence of favourable outcomes of higher levels of PA and lower levels of SB in both physical and mental health outcomes, and the emerging evidence that exercise can yield favourable COVID-19 outcomes, it is recommended that public health officials promote ways of increasing PA and reducing SB should further lockdowns occur, especially in populations with medical conditions that are improved by PA, such as type 1 and type 2 diabetes. Interventions designed for postlockdown should also consider that individuals may suffer from deconditioning during the lockdown period, especially in athletic populations and people with medical conditions.

References

- 1 World Health Organization. WHO coronavirus Disease (COVID-19) Dashboard. Available: <https://covid19.who.int> [Accessed 2 Jul 2020].
- 2 British Broadcasting Corporation. Coronavirus: the world in lockdown in maps and charts, 2020. Available: <https://www.bbc.co.uk/news/world-52103747> [Accessed 12 Aug 2020].
- 3 Hossain MM, Sultana A, Purohit N. Mental health outcomes of quarantine and isolation for infection prevention: a systematic umbrella review of the global evidence. *SSRN Journal*;19.
- 4 Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep* 1985;100:126.
- 5 Bazaco MC, Pereira MA, Wisniewski SR, et al. Is there a relationship between perceived neighborhood Contentedness and physical activity in young men and women. *J Urban Health* 2016;93:940–52.
- 6 Pedersen BK, Saltin B. Exercise as medicine - evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports* 2015;25(Suppl 3):1–72.
- 7 Schuch FB, Vancampfort D, Firth J, et al. Physical activity and incident depression: a meta-analysis of prospective cohort studies. *Am J Psychiatry* 2018;175:631–48.
- 8 Schuch FB, Stubbs B, Meyer J, et al. Physical activity protects from incident anxiety: a meta-analysis of prospective cohort studies. *Depress Anxiety* 2019;36:846–58.
- 9 Jacob L, Tully MA, Barnett Y. The relationship between physical activity and mental health in a sample of the UK public: a cross-sectional study during the implementation of COVID-19 social distancing measures. *Ment Health Phys Act* 2020;100345.
- 10 Schuch FB, Bulzing RA, Meyer J, et al. Associations of moderate to vigorous physical activity and sedentary behavior with depressive and anxiety symptoms in self-isolating people during the COVID-19 pandemic: a cross-sectional survey in Brazil. *Psychiatry Res*

2020;292:113339.

11 Simpson RJ, Katsanis E. The immunological case for staying active during the COVID-19 pandemic. *Brain Behav Immun* 2020;87:6–7.

12 Gallè F, Sabella EA, Da Molin G, et al. Understanding knowledge and behaviors related to CoViD–19 epidemic in Italian undergraduate students: the EPICO study. *Int J Environ Res Public Health* 2020;17:3481.

13 Tremblay MS, LeBlanc AG, Kho ME, et al. Systematic review of sedentary behaviour and health indicators in school-aged children and youth. *Int J Behav Nutr Phys Act* 2011;8:98.

14 Cunningham C, O' Sullivan R, Caserotti P, et al. Consequences of physical inactivity in older adults: a systematic review of reviews and meta-analyses. *Scand J Med Sci Sports* 2020;30:816–27.

15 Thorp AA, Owen N, Neuhaus M, et al. Sedentary behaviors and subsequent health outcomes in adults a systematic review of longitudinal studies, 1996-2011. *Am J Prev Med* 2011;41:207–15.

16 Meyer J, McDowell C, Lansing J. Changes in physical activity and sedentary behaviour due to the COVID-19 outbreak and associations with mental health in 3,052 US adults. *Cambridge Open Engage* 2020.

17 Sallis JF, Adlakha D, Oyeyemi A, et al. An international physical activity and public health research agenda to inform coronavirus disease-2019 policies and practices. *J Sport Health Sci* 2020;9:328–34.

18 Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med* 2009;6:e1000100.

- 19 Wells GA, Shea B, O'connell D. The Newcastle-Ottawa scale (NOS) for assessing the quality of nonrandomised studies in metaanalyses. Ottawa, ON: Ottawa Hospital Research Institute, 2009.
- 20 Modesti PA, Reboldi G, Cappuccio FP, et al. Panethnic differences in blood pressure in Europe: a systematic review and metaanalysis. *PLoS One* 2016;11:e0147601.
- 21 Ammar A, Chtourou H, Boukhris O, et al. COVID-19 home confinement negatively impacts social participation and life satisfaction: a worldwide multicenter study. *Int J Environ Res Public Health* 2020;17:6237.
- 22 Asiamah N, Opuni FF, Mends-Brew E, et al. Short-Term changes in behaviors resulting from COVID-19-Related social isolation and their influences on mental health in Ghana. *Community Ment Health J* 2021;57:79–92.
- 23 Assaloni R, Pellino VC, Puci MV, et al. Coronavirus disease (Covid-19): how does the exercise practice in active people with type 1 diabetes change? A preliminary survey. *Diabetes Res Clin Pract* 2020;166:108297.
- 24 Barrea L, Pugliese G, Framondi L, et al. Does Sars-Cov-2 threaten our dreams? Effect of quarantine on sleep quality and body mass index. *J Transl Med* 2020;18:318.
- 25 Barwais FA. Physical activity at home during the COVID-19 pandemic in the two Most-affected cities in Saudi Arabia. *Open Public Health J* 2020;13:470–6.
- 26 Biviá-Roig G, La Rosa VL, Gómez-Tébar M, et al. Analysis of the impact of the confinement resulting from covid-19 on the lifestyle and psychological wellbeing of Spanish pregnant women: an Internet-based cross-sectional survey. *Int J Environ Res Public Health* 2020;17:5933–14.
- 27 Bourdas DI, Zacharakis ED. Impact of COVID-19 Lockdown on physical activity in a sample of Greek adults. *Sports* 2020;8. doi:10.3390/sports8100139. [Epub ahead of print: 21 Oct 2020].

- 28 Bourdas DI, Zacharakis ED. Evolution of changes in physical activity over lockdown time: physical activity datasets of four independent adult sample groups corresponding to each of the last four of the six COVID-19 lockdown weeks in Greece. *Data Brief* 2020;32:106301.
- 29 Bowes A, Lomax L, Piasecki J. The impact of the COVID-19 lockdown on elite sportswomen. *Managing Sport and Leisure*;48:1–17.
- 30 Branley-Bell D, Talbot CV. Exploring the impact of the COVID-19 pandemic and UK lockdown on individuals with experience of eating disorders. *J Eat Disord* 2020;8:44.
- 31 Buoite Stella A, Ajčević M, Furlanis G, et al. Smart technology for physical activity and health assessment during COVID-19 lockdown. *J Sports Med Phys Fitness* 2020. doi:10.23736/S0022-4707.20.11373-2. [Epub ahead of print: 22 Oct 2020].
- 32 Callow DD, Arnold-Nedimala NA, Jordan LS, et al. The mental health benefits of physical activity in older adults survive the COVID-19 pandemic. *Am J Geriatr Psychiatry* 2020;28:1046–57.
- 33 Canello R, Soranna D, Zambra G, et al. Determinants of the lifestyle changes during COVID-19 pandemic in the residents of northern Italy. *Int J Environ Res Public Health* 2020;17:6287.
- 34 Caruso I, Di Molfetta S, Guarini F, et al. Reduction of hypoglycaemia, lifestyle modifications and psychological distress during lockdown following SARS-CoV-2 outbreak in type 1 diabetes. *Diabetes Metab Res Rev*;13.
- 35 Castañeda-Babarro A, Arbillaga-Etxarri A, Gutiérrez-Santamaría B, et al. Physical activity change during COVID-19 confinement. *Int J Environ Res Public Health* 2020;17:6878.
- 36 Castellini G, Cassioli E, Rossi E, et al. The impact of COVID -19 epidemic on eating disorders: A longitudinal observation of pre versus post psychopathological features in a sample of patients with eating disorders and a group of healthy controls. *Int J Eat Disord* 2020;53:1855–62.

- 37 Constandt B, Thibaut E, De Bosscher V, et al. Exercising in times of Lockdown: an analysis of the impact of COVID-19 on levels and patterns of exercise among adults in Belgium. *Int J Environ Res Public Health* 2020;17:4144.
- 38 Constant A, Conserve DF, Gallopel-Morvan K, et al. SocioCognitive factors associated with lifestyle changes in response to the COVID-19 epidemic in the general population: results from a cross-sectional study in France. *Front Psychol* 2020;11:579460.
- 39 Di Corrado D, Magnano P, Muzii B, et al. Effects of social distancing on psychological state and physical activity routines during the COVID-19 pandemic. *Sport Sci Health* 2020;16:619–24.
- 40 Di Stefano V, Battaglia G, Giustino V, et al. Significant reduction of physical activity in patients with neuromuscular disease during COVID-19 pandemic: the long-term consequences of quarantine. *J Neurol* 2021;268:20–6.
- 41 Đogaš Z, Lušić Kalcina L, Pavlinac Dodig I, et al. The effect of COVID-19 lockdown on lifestyle and mood in Croatian general population: a cross-sectional study. *Croat Med J* 2020;61:309–18.
- 42 Dutta K, Mukherjee R, Sen D. Effect of COVID-19 lockdown on sleep behavior and screen exposure time: an observational study among Indian school children. *Biol Rhythm Res* 2020;21:1–12.
- 43 Elran-Barak R, Mozeikov M. One month into the reinforcement of social distancing due to the COVID-19 outbreak: subjective health, health behaviors, and loneliness among people with chronic medical conditions. *Int J Environ Res Public Health* 2020;17:5403.
- 44 Endstrasser F, Braitto M, Linser M, et al. The negative impact of the COVID-19 lockdown on pain and physical function in patients with end-stage hip or knee osteoarthritis. *Knee Surg Sports Traumatol Arthrosc* 2020;28:2435–43.

- 45 Ernstsén L, Havnen A. Mental health and sleep disturbances in physically active adults during the COVID-19 lockdown in Norway: does change in physical activity level matter? *Sleep Med* 2020. doi:10.1016/j.sleep.2020.08.030. [Epub ahead of print: 05 Sep 2020].
- 46 Gallè F, Sabella EA, Ferracuti S, et al. Sedentary behaviors and physical activity of Italian undergraduate students during Lockdown at the time of CoViD-19 pandemic. *Int J Environ Res Public Health* 2020;17:6171–11.
- 47 Gallo LA, Gallo TF, Young SL, et al. The impact of isolation measures due to COVID-19 on energy intake and physical activity levels in Australian university students. *Nutrients* 2020;12:1865.
- 48 Gilic B, Ostojic L, Corluka M, et al. Contextualizing Parental/ Familial influence on physical activity in adolescents before and during COVID-19 pandemic: a prospective analysis. *Children* 2020;7:125–14.
- 49 Giustino V, Parroco AM, Gennaro A, et al. Physical activity levels and related energy expenditure during COVID-19 quarantine among the Sicilian active population: a cross-sectional online survey study. *Sustainability* 2020;12:4356.
- 50 He M, Xian Y, Lv X, et al. Changes in body weight, physical activity, and lifestyle during the Semi-lockdown period after the outbreak of COVID-19 in China: an online survey. *Disaster Med Public Health Prep* 2020:1–6.
- 51 Husain W, Ashkanani F. Does COVID-19 change dietary habits and lifestyle behaviours in Kuwait: a community-based cross-sectional study. *Environ Health Prev Med* 2020;25:61.
- 52 Ingram J, Maciejewski G, Hand CJ. Changes in diet, sleep, and physical activity are associated with differences in negative mood during COVID-19 Lockdown. *Front Psychol* 2020;11:588604.

- 53 Karuc J, Sorić M, Radman I, et al. Moderators of change in physical activity levels during restrictions due to COVID-19 pandemic in young urban adults. *Sustainability* 2020;12:6392.
- 54 Knell G, Robertson MC, Dooley EE, et al. Health behavior changes during covid-19 pandemic and subsequent “stay-at-home” orders. *Int J Environ Res Public Health* 2020;17:6268–16.
- 55 Kriaucioniene V, Bagdonaviciene L, Rodríguez-Pérez C, et al. Associations between changes in health behaviours and body weight during the covid-19 quarantine in Lithuania: the Lithuanian covidiet study. *Nutrients* 2020;12:3119–9.
- 56 López-Bueno R, Calatayud J, Andersen LL, et al. Immediate impact of the COVID-19 confinement on physical activity levels in Spanish adults. *Sustainability* 2020;12:5708.
- 57 López-Bueno R, Calatayud J, Casaña J, et al. COVID-19 confinement and health risk behaviors in Spain. *Front Psychol* 2020;11:1426.
- 58 Majumdar P, Biswas A, Sahu S. COVID-19 pandemic and lockdown: cause of sleep disruption, depression, somatic pain, and increased screen exposure of office workers and students of India. *Chronobiol Int* 2020;37:1191–200.
- 59 Mandelkorn U, Genzer S, Choshen-Hillel S, et al. Escalation of sleep disturbances amid the COVID-19 pandemic: a crosssectional International study. *J Clin Sleep Med* 2021;17:45–53.
- 60 Maugeri G, Castrogiovanni P, Battaglia G, et al. The impact of physical activity on psychological health during Covid-19 pandemic in Italy. *Heliyon* 2020;6:e04315.
- 61 Meyer J, McDowell C, Lansing J, et al. Changes in physical activity and sedentary behavior in response to COVID-19 and their associations with mental health in 3052 us adults. *Int J Environ Res Public Health* 2020;17:6469.
- 62 Mitra R, Moore SA, Gillespie M, et al. Healthy movement behaviours in children and youth during the COVID-19 pandemic: exploring the role of the neighbourhood environment. *Health Place* 2020;65:102418.

- 63 Mon-López D, de la Rubia Rianza A, Hontoria Galán M, et al. The impact of Covid-19 and the effect of psychological factors on training conditions of Handball players. *Int J Environ Res Public Health* 2020;17:6471.
- 64 Mon-López D, Bernardez-Vilaboa R, Fernandez-Balbuena AA, et al. The influence of COVID-19 isolation on physical activity habits and its relationship with convergence insufficiency. *Int J Environ Res Public Health* 2020;17:1–9.
- 65 Munasinghe S, Sperandei S, Freebairn L, et al. The impact of physical distancing policies during the COVID-19 pandemic on health and well-being among Australian adolescents. *J Adolesc Health* 2020;67:653–61.
- 66 Pellegrini M, Ponzio V, Rosato R, et al. Changes in Weight and Nutritional Habits in Adults with Obesity during the “Lockdown” Period Caused by the COVID-19 Virus Emergency. *Nutrients* 2020;12:2016.
- 67 Pillay L, Janse van Rensburg DCC, Jansen van Rensburg A, et al. Nowhere to hide: the significant impact of coronavirus disease 2019 (COVID-19) measures on elite and semi-elite South African athletes. *J Sci Med Sport* 2020;23:670–9.
- 68 Robinson E, Boyland E, Chisholm A, et al. Obesity, eating behavior and physical activity during COVID-19 lockdown: a study of UK adults. *Appetite* 2021;156:104853.
- 69 Rogers NT, Waterlow NR, Brindle H, et al. Behavioral change towards reduced intensity physical activity is disproportionately prevalent among adults with serious health issues or Self-Perception of high risk during the UK COVID-19 Lockdown. *Front Public Health* 2020;8:575091.
- 70 Romero-Blanco C, Rodríguez-Almagro J, Onieva-Zafra MD, et al. Physical activity and sedentary lifestyle in university students: changes during confinement due to the COVID-19 pandemic. *Int J Environ Res Public Health* 2020;17:6567.

- 71 Ruíz-Roso MB, de Carvalho Padilha P, Matilla-Escalante DC, et al. Changes of physical activity and Ultra-Processed food consumption in adolescents from different countries during Covid-19 pandemic: an observational study. *Nutrients* 2020;12:2289.
- 72 Ruiz-Roso MB, Knott-Torcal C, Matilla-Escalante DC, et al. COVID-19 Lockdown and changes of the dietary pattern and physical activity habits in a cohort of patients with type 2 diabetes mellitus. *Nutrients* 2020;12:2327.
- 73 Sánchez-Sánchez E, Ramírez-Vargas G, Avellaneda-López Y, et al. Eating habits and physical activity of the Spanish population during the covid-19 pandemic period. *Nutrients* 2020;12:2826–12.
- 74 Sankar P, Ahmed WN, Mariam Koshy V, et al. Effects of COVID-19 lockdown on type 2 diabetes, lifestyle and psychosocial health: a hospital-based cross-sectional survey from South India. *Diabetes Metab Syndr* 2020;14:1815–9.
- 75 Sañudo B, Fennell C, Sánchez-Oliver AJ. Objectively-assessed physical activity, sedentary behavior, smartphone use, and sleep patterns pre- and during-COVID-19 quarantine in young adults from Spain. *Sustainability* 2020;12:5890.
- 76 Savage MJ, James R, Magistro D, et al. Mental health and movement behaviour during the COVID-19 pandemic in UK university students: prospective cohort study. *Ment Health Phys Act* 2020;19:100357.
- 77 Schlichtiger J, Brunner S, Steffen J, et al. Mental health impairment triggered by the COVID-19 pandemic in a sample population of German students. *J Investig Med* 2020;68:1394–6.
- 78 Schlichtiger J, Steffen J, Huber BC, et al. Physical activity during COVID-19 lockdown in older adults. *J Sports Med Phys Fitness* 2020;61.
- 79 Srivastav AK, Sharma N, Samuel AJ. Impact of coronavirus disease-19 (COVID-19) lockdown on physical activity and energy expenditure among physiotherapy professionals and students

using web-based open E-survey sent through WhatsApp, Facebook and Instagram messengers. Clin Epidemiol Glob Health 2021;9:78–84.

80 Vetrovsky T, Frybova T, Gant I, et al. The detrimental effect of COVID-19 nationwide quarantine on accelerometer-assessed physical activity of heart failure patients. ESC Heart Fail 2020;7:2093–7.

81 Wang X, Lei SM, Le S, et al. Bidirectional influence of the COVID-19 pandemic Lockdowns on health behaviors and quality of life among Chinese adults. Int J Environ Res Public Health 2020;17. doi:10.3390/ijerph17155575. [Epub ahead of print: 02 08 2020].

82 Yang S, Guo B, Ao L, et al. Obesity and activity patterns before and during COVID-19 lockdown among Youths in China. Clin Obes 2020;10:e12416.

83 Yang Y, Koenigstorfer J. Determinants of physical activity maintenance during the Covid-19 pandemic: a focus on fitness apps. Transl Behav Med 2020;10:835–42.

84 Zenic N, Taiar R, Gilic B, et al. Levels and changes of physical activity in adolescents during the COVID-19 pandemic: Contextualizing urban vs. rural living environment. Appl Sci 2020;10:3997.

85 Muriel X, Courel-Ibáñez J, Cerezuela-Espejo V, et al. Training load and performance impairments in professional cyclists during COVID-19 Lockdown. Int J Sports Physiol Perform 2020;1:1–4.

86 Pietrobelli A, Pecoraro L, Ferruzzi A, et al. Effects of COVID-19 Lockdown on lifestyle behaviors in children with obesity living in Verona, Italy: a longitudinal study. Obesity 2020;28:1382–5.

87 American College of Sports Medicine. Staying physically active during the COVID-19 pandemic, 2020. Available: <https://www.acsm.org/read-research/newsroom/news->

[releases/news-detail/2020/03/16/staying-physically-active-during-covid-19-pandemic](https://www.acsm.org/read-research/newsroom/news-releases/news-detail/2020/03/16/staying-physically-active-during-covid-19-pandemic) [Accessed 16 Aug 2020].

88 World Health Organization. Stay physically active during selfquarantine, 2020. Available: <https://www.acsm.org/read-research/newsroom/news-releases/news-detail/2020/03/16/stayingphysically-active-during-covid-19-pandemic> [Accessed 16 Aug 2020].

89 Dwyer MJ, Pasini M, De Dominicis S, et al. Physical activity: benefits and challenges during the COVID-19 pandemic. *Scand J Med Sci Sports* 2020;30:1291–4.

90 Çolak TK, Acar G, Dereli EE, et al. Association between the physical activity level and the quality of life of patients with type 2 diabetes mellitus. *J Phys Ther Sci* 2015;28:142–7.

91 Mutlu EK, Mutlu C, Taskiran H, et al. Association of physical activity level with depression, anxiety, and quality of life in children with type 1 diabetes mellitus. *J Pediatr Endocrinol Metab* 2015;28:1273–8.

92 Åman J, Skinner TC, de Beaufort CE, et al. Associations between physical activity, sedentary behavior, and glycemic control in a large cohort of adolescents with type 1 diabetes: the Hvidoere Study Group on childhood diabetes. *Pediatr Diabetes* 2009;10:234–9.

93 MacMillan F, Kirk A, Mutrie N, et al. A systematic review of physical activity and sedentary behavior intervention studies in youth with type 1 diabetes: study characteristics, intervention design, and efficacy. *Pediatr Diabetes* 2014;15:175–89.

94 Trott M, Jackson SE, Firth J, et al. A comparative meta-analysis of the prevalence of exercise addiction in adults with and without indicated eating disorders. *Eat Weight Disord* 2020. doi:10.1007/s40519-019-00842-1. [Epub ahead of print: 01 Jan 2020].

95 Michie S, Atkins L, West R. The behaviour change wheel: a guide to designing interventions. Great Britain: Silverback Publishing, 2014.

- 96 UK Government. The health protection (coronavirus, business closure) (England) regulations 2020, 2020. Available: <https://www.legislation.gov.uk/uksi/2020/327/contents/made> [Accessed 16 Aug 2020].
- 97 Di Renzo L, Gualtieri P, Pivari F, et al. Eating habits and lifestyle changes during COVID-19 lockdown: an Italian survey. *J Transl Med* 2020;18:1–15.
- 98 Office for National Statistics. Coronavirus and homeworking in the UK: April 2020, 2020. Available: <https://www.ons.gov.uk/employmentandlabourmarket/peopleinwork/employmentandemployeetypes/bulletins/coronavirusandhomeworkingintheuk/latest> [Accessed 16 Aug 2020].
- 99 Sahlqvist S, Song Y, Ogilvie D. Is active travel associated with greater physical activity? The contribution of commuting and noncommuting active travel to total physical activity in adults. *Prev Med* 2012;55:206–11.
- 100 Foley L, Panter J, Heinen E, et al. Changes in active commuting and changes in physical activity in adults: a cohort study. *Int J Behav Nutr Phys Act* 2015;12:161.
- 101 Learnpatch. Research reveals impact of Covid-19 on working practices and remote working, 2020. Available: <https://learnpatch.com/2020/05/covid-remote-working/> [Accessed 16 Aug 2020].
- 102 Qin F, Song Y, Nassis GP, et al. Prevalence of insufficient physical activity, sedentary screen time and emotional well-being during the early days of the 2019 novel coronavirus (COVID-19) outbreak in China: a national cross-sectional study. *SSRN Journal* 2020.
- 103 Smith L, Jacob L, Trott M, et al. The association between screen time and mental health during COVID-19: a cross sectional study. *Psychiatry Res* 2020;292:113333.
- 104 de Haas M, Faber R, Hamersma M. How COVID-19 and the Dutch ‘intelligent lockdown’ change activities, work and travel behaviour: Evidence from longitudinal data in the Netherlands. *Transportation Research Interdisciplinary Perspectives* 2020;6:100150.

105 Ammar A, Brach M, Trabelsi K, et al. Effects of COVID-19 home confinement on eating behaviour and physical activity: results of the ECLB-COVID19 international online survey. *Nutrients* 2020;12:1583.

106 Monbiot G. The horror films got it wrong. This virus has turned us into caring neighbours, 2020. Available: <https://www.theguardian.com/commentisfree/2020/mar/31/virus-neighbours-covid-19> [Accessed 16 Aug 2020].

107 Demetriou C, Ozer BU, Essau CA. Self-report questionnaires. *Encyclopedia Clin Psychol* 2015:1–6.